



HAL
open science

Impact of domestic mould exposure on *Aspergillus* biomarkers and lung function in patients with chronic obstructive pulmonary disease

Emilie Fréalle, Gabriel Reboux, Olivier Le Rouzic, Nathalie Bautin, Marie-Capucine Willemin, Muriel Pichavant, Julie Delourme, Boualem Sendid, Philippe Gosset, Saad Nseir, et al.

► To cite this version:

Emilie Fréalle, Gabriel Reboux, Olivier Le Rouzic, Nathalie Bautin, Marie-Capucine Willemin, et al.. Impact of domestic mould exposure on *Aspergillus* biomarkers and lung function in patients with chronic obstructive pulmonary disease. *Environmental Research*, 2021, 195, pp.110850. 10.1016/j.envres.2021.110850 . hal-03537200

HAL Id: hal-03537200

<https://univ-fcomte.hal.science/hal-03537200>

Submitted on 14 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Impact of domestic mould exposure on *Aspergillus* biomarkers and lung**
2 **function in patients with chronic obstructive pulmonary disease**

3

4 Emilie Fréalle^{a,b*}, Gabriel Reboux^c, Olivier Le Rouzic^{b,d}, Nathalie Bautin^d, Marie-Capucine
5 Willemin^d, Muriel Pichavant^b, Julie Delourme^d, Boualem Sendid^{a,e}, Philippe Gosset^b, Saad
6 Nseir^f, Stéphanie Fry^d

7 ^aCHU Lille, Laboratoire de Parasitologie-Mycologie, F-59000 Lille, France

8 ^bUniv. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 – UMR9017 - CIIL -
9 Center for Infection and Immunity of Lille, F-59000 Lille, France

10 ^cChrono-Environnement UMR 6249 CNRS, Université de Bourgogne Franche-Comté &
11 Service de Parasitologie-Mycologie, CHU de Besançon, France

12 ^dCHU Lille, Clinique des Maladies Respiratoires, F-59000 Lille, France

13 ^eInserm U995, Université de Lille, France

14 ^fCHU Lille, Pôle de Réanimation, F-59000 Lille, France

15

16 *Corresponding author: Dr Emilie Fréalle, CHU de Lille, Service de Parasitologie-Mycologie,
17 Centre de Biologie Pathologie, Boulevard du Pr Jules Leclercq, 59037 Lille Cedex, France.

18 Tel: +33-3-20-44-55-77, Fax: +33-3-20-44-48-95, E-mail address: emilie.frealle@chru-lille.fr

19

20

21 **ABSTRACT**

22 Patients with chronic obstructive pulmonary disease (COPD) are frequently colonised or
23 sensitised by *Aspergillus*, but clinical significance remains unclear. Furthermore, little is
24 known on the impact of indoor mould exposure during COPD. In this study, we assessed the
25 relationship between domestic mould exposure, *Aspergillus* biomarkers and COPD severity
26 during acute exacerbation and at stable state. *Aspergillus* section *Fumigati* culture in sputum
27 and anti-*Aspergillus* antibodies detection (IgG and precipitins) were followed up in COPD
28 patients that were prospectively recruited during exacerbation (n=62), and underwent a visit at
29 stable state after 18 months (n=33). Clinical characteristics were collected at inclusion.
30 Electrostatic dust collectors (EDCs) were used to measure domestic mould contamination.
31 *Aspergillus* section *Fumigati* was more frequently detected during exacerbation (16.9%) than
32 at stable state (4.0%), but the frequency of patients presenting with anti-*Aspergillus* antibodies
33 was similar (32.2% and 33.3%, respectively). *Aspergillus* section *Fumigati* detection was
34 associated with a higher body-mass index (BMI) during exacerbation, whereas patients with
35 anti-*Aspergillus* antibodies presented a lower BMI and forced expiratory volume in 1 second,
36 as well as a higher frequency of inhaled corticoids and higher total mould and *Penicillium*
37 exposure at final visit (P<0.05). The frequency of patients with anti-*Aspergillus* antibodies
38 was higher for total mould counts >30 CFUs/cm² (P=0.03). Aspergillosis was diagnosed in 2
39 patients (6.1%) who presented increased levels of antibodies. Our data suggest that anti-
40 *Aspergillus* antibodies are associated with chronic lung function alteration and/or domestic
41 mould exposure, thereby supporting the consideration of indoor mould contamination and
42 anti-*Aspergillus* antibodies kinetics in COPD management.

43 **Keywords:** COPD, *Aspergillus*, fungi, environment, anti-*Aspergillus* antibodies

44

45 **Funding**

46 This work was supported by the Région Hauts-de-France (Environment Health
47 program, grant N°2010-12266 & 2010-12268), acting for a healthier population, and by
48 GILEAD Sciences, MSD, and Pfizer grants.

49

50

51

52 **1. Introduction**

53 Chronic obstructive pulmonary disease (COPD) is a chronic inflammatory disease,
54 which has a global mean prevalence of 12.4% in subjects aged ≥ 40 years in Europe (Blanco et
55 al., 2018), and is mainly caused by inhalation of cigarette smoke. The hallmark feature of this
56 disease is airflow limitation, but a substantial number of patients also suffer from acute
57 exacerbations (AECOPD), which are a major cause of morbidity, mortality and hospital
58 admission (Soler-Cataluña et al., 2005). AECOPD is mainly triggered by infections due to
59 bacteria or viruses (Domenech et al., 2013), which have been shown to be facilitated by
60 chronic inflammation and impaired lung immune defences (Le Rouzic et al., 2017; Pichavant
61 et al., 2015, 2014). In addition, COPD patients are also at risk for invasive pulmonary
62 aspergillosis (IPA). IPA has been increasingly reported with incidences varying from 1.6 to
63 3.9%, probably under-estimated, and is associated with a high mortality rate (72-100%)
64 (Delsuc et al., 2015; Guinea et al., 2010).

65 Increasing evidence supports the role of filamentous fungi in stable COPD. Indeed,
66 *Aspergillus* section *Fumigati* and anti-*A. fumigatus* antibodies are frequently detected in
67 COPD patients at stable state (Agarwal et al., 2010; Bafadhel et al., 2014). Although patients
68 with or without *Aspergillus* section *Fumigati*-positive culture show no difference in
69 exacerbation frequency or respiratory function, *Aspergillus* sensitisation was found to be
70 associated with a lower forced expiratory volume in 1 second (FEV1) (Bafadhel et al., 2014)
71 or bronchiectasis (Everaerts et al., 2017). Indoor mould exposure, which is involved in the
72 development and evolution of other respiratory diseases (Caillaud et al., 2018; French Agency
73 for Food, Environmental and Occupational Health & Safety (ANSES), 2016) could also play
74 a role in the pathophysiology of COPD. However, little data is available to assess this
75 possibility. One study evaluated fungal contamination using electrostatic dust fall collectors

76 (EDCs) and showed no difference of fungal exposure between COPD patients and controls
77 (Barrera et al., 2019). Nevertheless, another study has recently shown that indoor
78 environments were associated with greater COPD symptoms and lower FEV1 at stable state
79 (Tiew et al., 2020). The presence of indoor visible moulds was also identified as an
80 independent risk factor of persistent bronchial obstruction (Ciebiada et al., 2014), and
81 elevated indoor dust fungal DNA levels were found to be associated with low FEV1 or forced
82 vital capacity in asthmatic patients (Shendell et al., 2012). Furthermore, visible moulds were
83 found to be associated with FEV1 decrease in the general population (Devien et al., 2018).
84 Altogether, these data suggest mould exposure could contribute to airflow limitation during
85 COPD.

86 Here, our aim was to address the specific topic of mould exposure during COPD by
87 using EDCs, which are innovative tools used to perform quantitative and qualitative
88 characterisation of fungal flora (Ege et al., 2011; Dauchy et al., 2018; Viegas et al., 2019). We
89 assessed the relationship between domestic mould exposure measured with EDCs, *Aspergillus*
90 biomarkers (*Aspergillus* culture in sputum and anti-*Aspergillus* antibodies detection), and
91 COPD severity (lung function and frequency of exacerbation episodes) in patients who were
92 prospectively recruited during severe acute exacerbation requiring hospitalisation, and
93 underwent a final visit at stable state after 18 months.

94

95 **2. Methods**

96 *2.1. Patient recruitment and follow-up, data collection, clinical and environmental sampling*

97 Patients with acute exacerbation of COPD, who had no other pulmonary diseases
98 (tuberculosis, cancer, diffuse bronchiectasis, asthma, etc.) and had not received antifungal
99 systemic treatment within the 6 previous months, were prospectively recruited in the

100 Pneumology, Intensive Care or Post-Emergency services of Lille University Hospital (France)
101 from August 2011 to November 2015. Age, sex, GOLD stage, FEV1 at stable state, frequency
102 of exacerbation episodes in the 2 previous years, tobacco status, and corticoid treatments were
103 collected at inclusion. The results of the cytobacteriological examination of sputum and viral
104 analyses were also collected to determine the origin of exacerbation. Patients were asked to
105 attend a visit at stable state 18 months after inclusion.

106 Sputum and serum samples were collected at inclusion and after 18 months, at stable
107 state, for detection of *Aspergillus* section *Fumigati* and anti-*Aspergillus* antibodies. Home
108 mould exposure was assessed using EDCs that were exposed for 10 weeks in the patient's
109 bedroom.

110 The study was approved by the local ethics committee (North-West Ethics Committee,
111 University of Rouen, France, referral number 2010-031) and was registered at
112 ClinicalTrials.gov (NCT02318524).

113

114 2.2. *Mycological analysis of sputa and anti-Aspergillus antibodies detection*

115 Sputum samples were half diluted with a solution of acetylcysteine (Digest-EUR,
116 Eurobio), and 10 µL were cultured on half-diluted Sabouraud agar medium with 0.5 g/L
117 amikacin (SAB) at 30°C, chromogenic agar medium at 37°C and erythritol agar medium at
118 24°C for 7 days. Anti-*Aspergillus* IgG were detected using the Platelia™ *Aspergillus* IgG kit
119 (Bio-Rad, Marne-La-Coquette, France), and an in-house Ouchterlony double
120 immunodiffusion method was used for detection of anti-*A. fumigatus* precipitins. Patients
121 were considered to have anti-*Aspergillus* antibodies when levels over 9 UA/mL using ELISA,
122 or at least 2 precipitin arcs were detected.

123

124 2.3. Sampling and mycological analysis of indoor dust

125 Domestic fungal exposure was assessed from EDCs that were analysed by culture.
126 EDC washing consisted of shaking for 10 minutes in 20 mL of sterile NaCl 0.9% with 0.1%
127 Tween 80 in a Stomacher® 80 Biomaster (Life System), as previously described (Dauchy et
128 al., 2018; Scherer et al., 2014), and 60 µL of the collected solution were cultured on SAB at
129 30°C, dichloran - rose Bengal – chloramphenicol (DRBC) - benomyl and dichloran - glycerol
130 (DG18) agar media (Oxoid, Dardilly, France) at 24°C for 7 days. The choice of DG18
131 medium was based on the recommendation by the ISO 16000-17 norm for analysis of air
132 samples by culture (ISO 16000-17, 2008). The SAB medium and 30°C incubation were
133 selected for optimal detection thermophilic pathogenic moulds, especially *Aspergillus* spp.
134 The DRBC – benomyl medium was added for its inhibitory properties on microtubule growth
135 and hyphal elongation, which allows detection of slow-growing fungi (Jochová et al., 1993).
136 Moulds were identified by macroscopic and microscopic (morphology of conidiophores,
137 conidia and hyphae) characteristics (ANOFEL, 2018; de Hoog et al., 2001), and CFUs were
138 counted on the 3 media for each genus or species. When moulds were detected on several
139 culture media, the final number of colonies corresponded to the results from the medium with
140 the highest number of colonies for a given species/genus (ISO 16000-17, 2008).

141

142 2.4. Statistical analysis

143 Data are presented as mean±standard error mean. Comparison of FEV1 and mould
144 contamination levels in EDCs was assessed using the t-test or Mann–Whitney U-test in
145 GraphPad Prism 5 for parametric or nonparametric groups, respectively. For comparison of
146 three groups or more, the Kruskal–Wallis test was used, with Dunn’s multiple comparison test

147 as post-test to compare individual pairs of groups. The Fisher's exact test was used for
148 proportions. A p-value of <0.05 was considered statistically significant.

149

150 **3. Results**

151 *3.1. Patients' characteristics at inclusion and origin of exacerbations*

152 Sixty-two patients meeting inclusion criteria and no exclusion criteria were recruited.
153 The sex ratio was 2.6 (46 men and 17 women), and the mean age was 63.3±10.4 years-old.
154 Nine patients had mild (GOLD II stage), 28 had severe (stage III) and 25 had very severe
155 COPD (stage IV). Overall, the mean FEV1 reached 35.3±2.0% of predicted and the mean
156 body-mass index (BMI) was 25.2±0.8 kg/m². All the patients had a previous history of
157 smoking. Mean pack years was 63.4±4.2, whereas 17/62 patients (27.4%) were current
158 smokers. In the 2 previous years, 33% presented no exacerbation episode, whereas 24%
159 presented at least 3 episodes. Inhaled corticoids were used as maintenance treatment in 80.6%
160 of patients. During exacerbation, a bacterial infection was confirmed for 21 patients (33.9%),
161 and among them the most frequent bacteria were *Streptococcus pneumoniae* and *Haemophilus*
162 *influenzae* (both isolated in 6 sputa, i.e. 28.6% of patients with documented bacterial
163 exacerbation). *Pseudomonas aeruginosa* was isolated in 5 patients (23.8%), and *Moraxella*
164 *catarrhalis* in 3 patients (14.3%). Five patients (8.1%) presented a viral infection.

165

166 *3.2. Frequency and clinical significance of Aspergillus positive biomarkers during* 167 *exacerbation and at stable state*

168 At inclusion, *Aspergillus* section *Fumigati* was detected in 10 out of the 59 patients
169 who were able to expectorate (16.9%). Twenty patients (32.2%) presented anti-*Aspergillus*
170 antibodies, with IgG levels varying from 2 to over 80 UA/mL and 0 to 8 anti-*A. fumigatus*

171 precipitins. Five patients had both positive culture and anti-*Aspergillus* antibodies (8.1%). No
172 patient presented current *Aspergillus* infection, but 3 patients (N°45, N°52 and N°60) had a
173 history of *Aspergillus* infection 3 to 6 years before inclusion. Patients N°52 and N°60, but not
174 N°45, received voriconazole. Mortality reached 19.4% during the 18-month follow-up period.

175 Thirty-three patients underwent the final visit at stable state. Sputa were sampled for
176 25 out of 33 patients, and *Aspergillus* section *Fumigati* was detected in 1/25 patients (4.0%),
177 yielding a lower frequency at stable state than at inclusion, but this difference was not
178 significant. The frequency of patients with anti-*Aspergillus* antibodies reached 33.3%, being
179 similar at stable state and at inclusion. Only one patient had both positive culture and anti-
180 *Aspergillus* antibodies (3.0%). When comparing the individual evolution of *Aspergillus*
181 biomarkers, the 4 patients with positive culture became negative at stable state, the only
182 patient who had positive culture at final visit was negative at inclusion, and 20 negative
183 patients remained negative (Fig. 1). Anti-*Aspergillus* antibodies were detected in 3 out of 23
184 patients who were negative at inclusion. Among the 10 patients with anti-*Aspergillus*
185 antibodies at inclusion, 8 remained positive, 3 having stable antibody levels and 5 increased
186 antibody levels at the final visit. Aspergillosis was diagnosed at final visit in 2 patients
187 (6.1%), who had increased anti-*Aspergillus* antibodies (initial diagnosis in patient N°10 and
188 recurrence in patient N°52). Both patients presented clinical (worsening respiratory
189 insufficiency in spite of appropriate antibiotic therapy and ventilatory support), and
190 radiological features of aspergillosis (abnormal medical imaging by CT scan). But only
191 patient N°10 had positive *Aspergillus* section *Fumigati* sputum culture. For patient N°10,
192 evolution was favorable after a 6 month treatment with voriconazole. A decrease in anti-
193 *Aspergillus* antibodies was observed after the end of the treatment (Fig. 2). Patient N°52 was
194 initially treated with voriconazole, which was replaced with liposomal amphotericin B after 1

195 month. However, he died after 20 days. No aspergillosis was diagnosed in any of the other
196 patients, even when significant levels of anti-*Aspergillus* antibodies were detected.

197

198 3.3. Relationship between *Aspergillus* biomarkers and patients characteristics at inclusion

199 Mean FEV1 tended to be lower in patients with *Aspergillus* section *Fumigati* positive
200 culture (P=0.06; Table 1, Fig. 3A) than without, and was significantly lower in patients with
201 (P=0.01; Table 1, Fig. 3B) than without anti-*Aspergillus* antibodies. Frequency of inhaled
202 corticoid treatment was significantly higher in patients than in patients with than without anti-
203 *Aspergillus* antibodies. Mean BMI was significantly higher in patients with *Aspergillus*
204 section *Fumigati* positive culture, and lower in patients with anti-*Aspergillus* antibodies.
205 Smoking status and frequency of exacerbation with confirmed bacterial origin were similar in
206 patients with or without anti-*Aspergillus* antibodies, or in patients with positive or negative
207 culture (Table 1).

208

209 3.4. Assessment of domestic mould contamination and relationship with *Aspergillus* 210 biomarkers at inclusion and final visit

211 EDCs were sent to the laboratory by 51 out of 62 patients. Assessment of domestic
212 mould contamination yielded 0.6 to 98.1 CFUs/cm², representing 1 to 13 genera or species.
213 Mould frequencies and counts were higher on DG18 for most genera and species, except for
214 *Aspergillus* section *Fumigati* and mucorales which were more frequently detected on SAB
215 and Benomyl, respectively (Supplementary Table). Overall counts resulting from the 3 media
216 showed that *Penicillium* spp., *Cladosporium* spp., *Aspergillus* section *Fumigati*, *Aspergillus*
217 series *Versicolores* and Mucorales were the most frequent moulds (in 90.2%, 64.7%, 49%,
218 39.2% and 33% of EDCs, respectively; Table 2). *Aspergillus* section *Nigri*, *Fusarium* spp. and

219 *Scopulariopsis* spp., which are thermotolerant pathogenic moulds, were also frequently
220 isolated (17.6-25.5%). *Aspergillus* genera represented 1.5-92.5% of total fungal flora (median
221 22.2%), with *Aspergillus* section *Fumigati* corresponding to 0-100% of *Aspergillus* spp.
222 (median 16.7%) (Supplementary Fig.1). Overall mould contamination level and *Penicillium*
223 spp. concentrations tended to be higher in patients with positive vs. negative culture or
224 detection of anti-*Aspergillus* antibodies at inclusion (Fig. 4A-B, 4D-E), and were significantly
225 higher in patients with positive vs. negative detection of anti-*Aspergillus* antibodies at the
226 final visit ($P<0.05$; Fig. 4C, 4F). Furthermore, when we analysed the repartition of total
227 mould contamination level, we identified a group of 8 dwellings (15.7%) presenting a higher
228 contamination level, >30 CFUs/cm² (Supplementary Fig. 2). When comparing the frequency
229 of patients with or without anti-*Aspergillus* antibodies for dwellings with total contamination
230 level of \leq or >30 CFUs/cm², we found that the frequency of patients with anti-*Aspergillus*
231 antibodies tended to be higher at inclusion and was significantly higher at final visit in
232 dwellings with contamination levels >30 CFUs/cm² ($P=0.03$; Table 3). Since *Aspergillus*
233 section *Fumigati* was only detected in a single patient's sputum at the final visit, the impact of
234 domestic mould contamination on this biomarker could not be assessed. We found no
235 significant difference in mould contamination level in patients with positive culture or anti-
236 *Aspergillus* antibodies when *Aspergillus* section *Fumigati*, *Aspergillus* spp., or *Cladosporium*
237 spp. concentrations were compared (Supplementary Fig. 3). No threshold yielding significant
238 differences between the frequencies of patients with positive or negative culture or detection
239 of anti-*Aspergillus* antibodies could be identified for these specific moulds or for *Penicillium*
240 spp.

241

242 *3.5. Relationship between domestic mould contamination and COPD stage or exacerbation*
243 *frequency*

244 Total mould counts tended to be lower in stage-II patients than in stage-III or -IV
245 patients, and *Aspergillus* section *Fumigati* counts tended to be higher in stage-IV than in
246 stage- II or -III patients (Fig. 5A), but these differences were not significant. *Aspergillus* spp.,
247 *Penicillium* spp. and *Cladosporium* spp. counts were similar in stage-II, -III and -IV patients
248 (Supplementary Fig. 4A). Total mould counts, *Aspergillus* section *Fumigati*, *Aspergillus* spp.,
249 *Penicillium* spp. and *Cladosporium* spp. counts were also similar in patients with 0, 1, 2 or
250 more than 3 exacerbations during the previous 2 years (Fig. 5B, Supplementary Fig. 4B).

251

252 **4. Discussion**

253 Our study showed a 16.9% frequency of *Aspergillus* section *Fumigati* detection in
254 sputum during AECOPD and 4.0% at stable state. These frequencies were lower than those
255 reported previously (28%-37%) (Bafadhel et al., 2014). However, lower quantities of sputum
256 were used in our study. A limitation could also be the absence of repeated samples to detect
257 *Aspergillus* section *Fumigati* colonisation in our study. The higher frequency of *Aspergillus*
258 section *Fumigati* we found at inclusion is consistent with impaired immune response during
259 exacerbation (Berenson et al., 2014), which could facilitate *Aspergillus* colonisation.

260 Conversely, the frequencies of patients with anti-*Aspergillus* antibodies were similar
261 during exacerbation (32.2%) and at stable state (33.3%). Both frequencies were higher than
262 those reported in previous studies which used skin prick test and/or *A. fumigatus* IgE or IgG
263 detection by ImmunoCAP (8.5-18% in Agarwal et al., 2010; Bafadhel et al., 2014; Everaerts
264 et al., 2017), whereas IgG detection by ELISA and precipitins were used in our study. These
265 similar frequencies during exacerbation and at stable state suggest that anti-*Aspergillus*
266 antibodies are more related to chronic lung-function alteration and/or environmental
267 exposures than acute disorders during COPD. This hypothesis is also supported by the

268 association of anti-*Aspergillus* antibodies with a lower FEV1, and the higher frequency of
269 inhaled corticoids in patients with anti-*Aspergillus* antibodies. The lower BMI in patients with
270 anti-*Aspergillus* antibodies is consistent with the association of a lower BMI with chronic
271 *Aspergillus* infections (Ohara et al., 2016). Lower BMI was also identified among the risk
272 factors of pneumonia during COPD, together with lower FEV1 or higher inhaled
273 corticosteroid dose (Agusti et al., 2018; Crim et al., 2009). The higher BMI in patients with
274 positive detection of *Aspergillus* section *Fumigati* is surprising. Since *A. fumigatus*
275 colonisation in COPD patients with acute exacerbation is a risk factor for invasive
276 aspergillosis (Blot et al., 2012; Guinea et al., 2010; Khasawneh et al., 2006), this result could
277 be related to a higher risk of developing invasive aspergillosis during exacerbation for patients
278 with higher BMI.

279 Anti-*Aspergillus* antibody detection was used as a diagnostic criterion for the 2
280 patients who developed aspergillosis in our study. Although a positive *A. fumigatus* serum test
281 has been proposed as a criterion for IPA diagnosis (Bulpa et al., 2007), the clinical
282 significance of anti-*Aspergillus* antibodies during COPD remains unclear, and this marker is
283 not among the criteria that were proposed by Blot et al. for critically ill patients (Blot et al.,
284 2012). In our study, 3 other patients presented increased anti-*Aspergillus* antibodies or
285 became positive, and were not diagnosed with *Aspergillus* infection in the absence of clinical
286 and radiological criteria for aspergillosis. Since elevated IgG against *A. fumigatus* are usually
287 considered to indicate *Aspergillus* infection (Uffredi et al., 2003), these data underscore the
288 need for further studies on the kinetics of *Aspergillus* IgG and precipitins during COPD to
289 evaluate its clinical significance and usefulness for the diagnosis of acute infections.

290 The use of EDCs for assessment of mould contamination yielded the first data on
291 domestic fungal contamination in our region, showing frequencies of *Penicillium*,
292 *Cladosporium*, and *Aspergillus* series *Versicolores* that were similar to those reported in other

293 studies in France (Dallongeville et al., 2015; Reboux et al., 2009). However, higher
294 frequencies of pathogenic thermophilic moulds, such as *Aspergillus* section *Fumigati* and
295 Mucorales were observed in our study. Our data confirm that this easy-to-use method, which,
296 contrary to air sampling, does not require specific equipment, and is representative of fungal
297 exposure over several weeks, is a promising alternative for quantitative measurement of
298 mould contamination in dwellings (Fréalles et al., 2017).

299 An interesting finding in our study was the higher domestic mould exposure (total
300 counts and *Penicillium* spp. concentrations) of patients with anti-*Aspergillus* antibodies,
301 which supports the idea that mould exposure is a risk factor for development of anti-
302 *Aspergillus* antibodies during COPD. The association between anti-*Aspergillus* antibodies and
303 increased *Penicillium* (but not other mould concentrations) could be related to the
304 predominance of this fungal genus in indoor environments, and to increased indoor
305 concentrations of *Penicillium* in dwellings with indoor air-quality problems (Cooley et al.,
306 1998). The tendency to a higher total mould contamination level in dwellings from stage-III
307 and stage-IV patients and *Aspergillus* section *Fumigati* counts in stage-IV patients is
308 consistent with previous reports showing association between mould exposure and airway
309 obstruction in asthmatic patients or healthy subjects (Ciebiada et al., 2014; Devien et al.,
310 2018; Shendell et al., 2012), and supports the recent finding of an association between the
311 abundance of fungal allergens in indoor environments and greater COPD symptoms or lower
312 FEV1 (Tiew et al., 2020). Beyond the significant association between anti-*Aspergillus*
313 antibodies and total fungal counts in EDCs, the higher frequency of patients with anti-
314 *Aspergillus* antibodies in dwellings with total mould contamination levels of >30 CFUs/cm²
315 suggests that this threshold could be used to define abnormal indoor mould exposure, as well
316 as the need for patient's home assessment by a professional (such as a medical indoor-
317 environment counsellor) and remediation measures. Since the lack of threshold values is a

318 major problem for assessment of mould exposure, and there are currently no guidelines on
319 acceptable levels of fungal exposure in dust, this definition of a threshold for EDCs represents
320 an important finding for the further use of this method in practical assessment of dwellings.

321 The absence of association between *Aspergillus* section *Fumigati* in EDCs and
322 *Aspergillus* section *Fumigati* detection in sputum or antibody detection could be due to the
323 fact that, although high concentrations of airborne *Aspergillus* section *Fumigati* can be found
324 in dwellings (Paluch et al., 2019) or near specific reservoirs such as compost (Williams et al.,
325 2019), this species is usually not predominant in the environment (Reboux et al., 2009;
326 Vesper et al., 2007). The establishment of a relationship between *Aspergillus* section *Fumigati*
327 exposure and colonisation/anti-*Aspergillus* antibodies could also be biased by the
328 pathogenicity of these thermophilic species, which can yield transient colonisation, making
329 low doses have a higher effect than exposure to high concentrations of non-pathogenic species
330 (Eduard, 2009). The fact that differences were found only when total counts or *Penicillium*
331 spp. concentrations were compared (according to anti-*Aspergillus* antibodies status)
332 underscores the difficulty of assessing the pathogenic effect of moulds which are usually
333 present in low concentrations in dwellings. Using total fungal counts, which mainly reflect the
334 usual predominant species (i.e. *Penicillium* and *Cladosporium*), as a global indicator of fungal
335 risk could allow us to indirectly assess the risk related to minority species (such as *Aspergillus*
336 section *Fumigati*) or to species poorly detected by culture (such as *Stachybotrys chartarum*).
337 The use of real-time PCR for DNA quantification of these species could also be helpful in
338 further studies.

339 Our study was limited by the quite low number of patients. Nevertheless, this number
340 is close to the one that was reported in several other studies where COPD patients were
341 included during exacerbation (which is more limiting than inclusion at stable state), and
342 sputum was analysed for bacterial and fungal culture (Bafadhel et al., 2014). Home evaluation

343 with EDCs was a supplementary cause of refusal and further limited the recruitment. Despite
344 this quite low number of patients, we were able to confirm data that were observed in other
345 studies (such as a lower FEV1 or higher frequency of inhaled corticoids in patients with anti-
346 *Aspergillus* antibodies (Bafadhel et al., 2014; Everaerts et al., 2017)), and provided new data
347 on the relationship between mould exposure and bioclinical parameters during COPD.
348 However, having a greater number of patients would have been helpful to confirm some
349 tendencies that were observed (e.g., relationship between mould exposure and COPD stage),
350 or to assess more reliably the relationship between mould exposure and exacerbation
351 frequency.

352 Despite this limitation, our data suggest that *Aspergillus* colonisation could be
353 facilitated by impaired immune response during COPD exacerbation, whereas anti-
354 *Aspergillus* antibodies are rather associated with chronic lung-function alteration. The
355 association of mould exposure with anti-*Aspergillus* antibodies and association of anti-
356 *Aspergillus* antibodies with a lower FEV1 suggest abnormal indoor mould contamination
357 could contribute to the alteration of lung function during COPD, thereby supporting the
358 consideration of domestic mould exposure in the management of COPD patients. Further
359 studies are needed to assess the clinical significance of anti-*Aspergillus* antibody kinetics and
360 the potential role of mould exposure in the development of aspergillosis during COPD.

361

362 **Declaration of competing interest**

363 E. Fréalle reports research grants from Gilead, MSD and Pfizer, and travel grants for
364 congress and conference attendance by Gilead. The other authors have no conflicts of interest
365 to disclose with the submitted work.

366

367 **Credit author statement**

368 EF, GR, PG and SF conceived the study and designed the project. OLR, NB, MCW,
369 JD, SN and SF recruited the patients and collected the data. EF, GR and SF analysed the data.
370 EF wrote the manuscript draft. All authors critically revised the manuscript and approved the
371 submitted version.

372

373 **Acknowledgements**

374 We would like to dedicate this work to the memory of Eduardo Dei-Cas and Isabelle
375 Tillie-Leblond, who brought their expertise in fungal infections and respiratory diseases for
376 the initial study design. We are grateful to the Center for Clinical Investigation (CIC) of Lille
377 University Hospital, to Stéphanie Delvart, Sandrine Bonaventure, Dorothée Monvillers, and
378 Laurence Dumortier for valuable technical assistance, and all the physicians who participated
379 to the recruitment and follow-up of patients. This work was partially presented as an oral
380 communication at the TIMM-8 in 2017, at the congress of the “French Medical Mycology
381 Society” (SFMM) in 2017 and 2019, and at the congress of the “French Society of
382 Environmental Health” (SFSE) in 2018.

383

384

385

386 **References**

- 387 Agarwal, R., Hazarika, B., Gupta, D., Aggarwal, A.N., Chakrabarti, A., Jindal, S.K., 2010.
388 *Aspergillus* hypersensitivity in patients with chronic obstructive pulmonary disease:
389 COPD as a risk factor for ABPA? *Med. Mycol.* 48, 988–994.
390 <https://doi.org/10.3109/13693781003743148>
- 391 Agusti, A., Fabbri, L.M., Singh, D., Vestbo, J., Celli, B., Franssen, F.M.E., Rabe, K.F., Papi,
392 A., 2018. Inhaled corticosteroids in COPD: friend or foe? *Eur. Respir. J.* 52.
393 <https://doi.org/10.1183/13993003.01219-2018>
- 394 ANOFEL, 2018. *Parasitologie et mycologie médicales - guide des analyses et méthodes*,
395 Elsevier. ed.
- 396 Bafadhel, M., McKenna, S., Agbetile, J., Fairs, A., Desai, D., Mistry, V., Morley, J.P.,
397 Pancholi, M., Pavord, I.D., Wardlaw, A.J., Pashley, C.H., Brightling, C.E., 2014.
398 *Aspergillus fumigatus* during stable state and exacerbations of COPD. *Eur. Respir. J.*
399 43, 64–71. <https://doi.org/10.1183/09031936.00162912>
- 400 Barrera, C., Rocchi, S., Degano, B., Soumagne, T., Laurent, L., Bellanger, A.-P., Laplante, J.-
401 J., Millon, L., Dalphin, J.-C., Reboux, G., 2019. Microbial exposure to dairy farmers’
402 dwellings and COPD occurrence. *Int. J. Environ. Health Res.* 29, 387–399.
403 <https://doi.org/10.1080/09603123.2018.1545900>
- 404 Berenson, C.S., Kruzel, R.L., Eberhardt, E., Dolnick, R., Minderman, H., Wallace, P.K.,
405 Sethi, S., 2014. Impaired innate immune alveolar macrophage response and the
406 predilection for COPD exacerbations. *Thorax* 69, 811–818.
407 <https://doi.org/10.1136/thoraxjnl-2013-203669>
- 408 Blanco, I., Diego, I., Bueno, P., Fernández, E., Casas-Maldonado, F., Esquinas, C., Soriano,
409 J.B., Miravittles, M., 2018. Geographical distribution of COPD prevalence in Europe,

410 estimated by an inverse distance weighting interpolation technique. *Int. J. Chron.*
411 *Obstruct. Pulmon. Dis.* 13, 57–67. <https://doi.org/10.2147/COPD.S150853>

412 Blot, S.I., Taccone, F.S., Van den Abeele, A.-M., Bulpa, P., Meersseman, W., Brusselaers, N.,
413 Dimopoulos, G., Paiva, J.A., Misset, B., Rello, J., Vandewoude, K., Vogelaers, D.,
414 AspICU Study Investigators, 2012. A clinical algorithm to diagnose invasive
415 pulmonary aspergillosis in critically ill patients. *Am. J. Respir. Crit. Care Med.* 186,
416 56–64. <https://doi.org/10.1164/rccm.201111-1978OC>

417 Bulpa, P., Dive, A., Sibille, Y., 2007. Invasive pulmonary aspergillosis in patients with
418 chronic obstructive pulmonary disease. *Eur. Respir. J.* 30, 782–800.
419 <https://doi.org/10.1183/09031936.00062206>

420 Caillaud, D., Leynaert, B., Keirsbulck, M., Nadif, R., mould ANSES working group, 2018.
421 Indoor mould exposure, asthma and rhinitis: findings from systematic reviews and
422 recent longitudinal studies. *Eur. Respir. Rev. Off. J. Eur. Respir. Soc.* 27.
423 <https://doi.org/10.1183/16000617.0137-2017>

424 Ciebiada, M., Domagała, M., Gorska-Ciebiada, M., Gorski, P., 2014. Risk factors associated
425 with irreversible airway obstruction in nonsmoking adult patients with severe asthma.
426 *Allergy Asthma Proc.* 35, 72–79. <https://doi.org/10.2500/aap.2014.35.3785>

427 Cooley, J.D., Wong, W.C., Jumper, C.A., Straus, D.C., 1998. Correlation between the
428 prevalence of certain fungi and sick building syndrome. *Occup. Environ. Med.* 55,
429 579–584. <https://doi.org/10.1136/oem.55.9.579>

430 Crim, C., Calverley, P.M.A., Anderson, J.A., Celli, B., Ferguson, G.T., Jenkins, C., Jones,
431 P.W., Willits, L.R., Yates, J.C., Vestbo, J., 2009. Pneumonia risk in COPD patients
432 receiving inhaled corticosteroids alone or in combination: TORCH study results. *Eur.*
433 *Respir. J.* 34, 641–647. <https://doi.org/10.1183/09031936.00193908>

434 Dallongeville, A., Le Cann, P., Zmirou-Navier, D., Chevrier, C., Costet, N., Annesi-Maesano,
435 I., Blanchard, O., 2015. Concentration and determinants of molds and allergens in
436 indoor air and house dust of French dwellings. *Sci. Total Environ.* 536, 964–972.
437 <https://doi.org/10.1016/j.scitotenv.2015.06.039>

438 Dauchy, C., Bautin, N., Nseir, S., Reboux, G., Wintjens, R., Le Rouzic, O., Sendid, B.,
439 Viscogliosi, E., Le Pape, P., Arendrup, M.C., Gosset, P., Fry, S., Fréalle, E., 2018.
440 Emergence of *Aspergillus fumigatus* azole resistance in azole-naïve patients with
441 chronic obstructive pulmonary disease and their homes. *Indoor Air* 28, 298–306.
442 <https://doi.org/10.1111/ina.12436>

443 de Hoog, G.S., Guarro, J., Gene, J., Figueras, M., 2001. *Atlas of Clinical Fungi*, 2nd ed. Amer
444 Society for Microbiology.

445 Delsuc, C., Cottereau, A., Frealle, E., Bienvenu, A.-L., Dessein, R., Jarraud, S., Dumitrescu,
446 O., Le Maréchal, M., Wallet, F., Friggeri, A., Argaud, L., Rimmelé, T., Nseir, S.,
447 Ader, F., 2015. Putative invasive pulmonary aspergillosis in critically ill patients with
448 chronic obstructive pulmonary disease: a matched cohort study. *Crit. Care* 19, 421.
449 <https://doi.org/10.1186/s13054-015-1140-1>

450 Devien, L., Giovannelli, J., Cuny, D., Matran, R., Amouyel, P., Hulo, S., Edmé, J.L.,
451 Dauchet, L., 2018. Sources of household air pollution: The association with lung
452 function and respiratory symptoms in middle-aged adult. *Environ. Res.* 164, 140–148.
453 <https://doi.org/10.1016/j.envres.2018.02.016>

454 Domenech, A., Puig, C., Martí, S., Santos, S., Fernández, A., Calatayud, L., Dorca, J.,
455 Ardanuy, C., Liñares, J., 2013. Infectious etiology of acute exacerbations in severe
456 COPD patients. *J. Infect.* 67, 516–523. <https://doi.org/10.1016/j.jinf.2013.09.003>

457 Eduard, W., 2009. Fungal spores: a critical review of the toxicological and epidemiological
458 evidence as a basis for occupational exposure limit setting. *Crit. Rev. Toxicol.* 39,
459 799–864. <https://doi.org/10.3109/10408440903307333>

460 Ege, M.J., Mayer, M., Normand, A.-C., Genuneit, J., Cookson, W.O.C.M., Braun-Fahrländer,
461 C., Heederik, D., Piarroux, R., von Mutius, E., GABRIELA Transregio 22 Study
462 Group, 2011. Exposure to environmental microorganisms and childhood asthma. *N.*
463 *Engl. J. Med.* 364, 701–709. <https://doi.org/10.1056/NEJMoa1007302>

464 Everaerts, S., Lagrou, K., Dubbeldam, A., Lorent, N., Vermeersch, K., Van Hoeyveld, E.,
465 Bossuyt, X., Dupont, L.J., Vanaudenaerde, B.M., Janssens, W., 2017. Sensitization to
466 *Aspergillus fumigatus* as a risk factor for bronchiectasis in COPD. *Int. J. Chron.*
467 *Obstruct. Pulmon. Dis.* 12, 2629–2638. <https://doi.org/10.2147/COPD.S141695>

468 Fréalle, E., Bex, V., Reboux, G., Roussel, S., Bretagne, S., 2017. Classical and molecular
469 methods for identification and quantification of domestic moulds. *Rev. Mal. Respir.*
470 34, 1124–1137. <https://doi.org/10.1016/j.rmr.2017.01.009>

471 French Agency for Food, Environmental and Occupational Health & Safety (ANSES), 2016.
472 ANSES opinion and report on mould in buildings.

473 Guinea, J., Torres-Narbona, M., Gijón, P., Muñoz, P., Pozo, F., Peláez, T., de Miguel, J.,
474 Bouza, E., 2010. Pulmonary aspergillosis in patients with chronic obstructive
475 pulmonary disease: incidence, risk factors, and outcome. *Clin. Microbiol. Infect.* 16,
476 870–877. <https://doi.org/10.1111/j.1469-0691.2009.03015.x>

477 ISO 16000-17, 2008. Indoor air — Part 17: Detection and enumeration of moulds — Culture-
478 based method.

479 Jochová, J., Rupeš, I., Peberdy, J.F., 1993. Effect of the microtubule inhibitor benomyl on
480 protein secretion in *Aspergillus nidulans*. *Mycol. Res.* 97, 23–27.
481 [https://doi.org/10.1016/S0953-7562\(09\)81107-6](https://doi.org/10.1016/S0953-7562(09)81107-6)

482 Khasawneh, F., Mohamad, T., Moughrabieh, M.K., Lai, Z., Ager, J., Soubani, A.O., 2006.
483 Isolation of *Aspergillus* in critically ill patients: a potential marker of poor outcome. J.
484 Crit. Care 21, 322–327. <https://doi.org/10.1016/j.jccrc.2006.03.006>

485 Le Rouzic, O., Pichavant, M., Frealle, E., Guillon, A., Si-Tahar, M., Gosset, P., 2017. Th17
486 cytokines: novel potential therapeutic targets for COPD pathogenesis and
487 exacerbations. Eur. Respir. J. 50. <https://doi.org/10.1183/13993003.02434-2016>

488 Ohara, S., Tazawa, Y., Tanai, C., Tanaka, Y., Noda, H., Horiuchi, H., Usui, K., 2016. Clinical
489 characteristics of patients with *Aspergillus* species isolation from respiratory samples:
490 Comparison of chronic pulmonary aspergillosis and colonization. Respir. Investig. 54,
491 92–97. <https://doi.org/10.1016/j.resinv.2015.08.007>

492 Paluch, M., Lejeune, S., Hecquet, E., Prévotat, A., Deschildre, A., Fréalle, E., 2019. High
493 airborne level of *Aspergillus fumigatus* and presence of azole-resistant TR34/L98H
494 isolates in the home of a cystic fibrosis patient harbouring chronic colonisation with
495 azole-resistant H285Y *A. fumigatus*. J. Cyst. Fibros. 18, 364–367.
496 <https://doi.org/10.1016/j.jcf.2019.01.001>

497 Pichavant, M., Rémy, G., Bekaert, S., Le Rouzic, O., Kervoaze, G., Vilain, E., Just, N., Tillie-
498 Leblond, I., Trottein, F., Cataldo, D., Gosset, P., 2014. Oxidative stress-mediated
499 iNKT-cell activation is involved in COPD pathogenesis. Mucosal Immunol. 7, 568–
500 578. <https://doi.org/10.1038/mi.2013.75>

501 Pichavant, M., Sharan, R., Le Rouzic, O., Olivier, C., Hennegrave, F., Rémy, G., Pérez-Cruz,
502 M., Koné, B., Gosset, Pierre, Just, N., Gosset, Philippe, 2015. IL-22 Defect During
503 *Streptococcus pneumoniae* Infection Triggers Exacerbation of Chronic Obstructive
504 Pulmonary Disease. EBioMedicine 2, 1686–1696.
505 <https://doi.org/10.1016/j.ebiom.2015.09.040>

506 Reboux, G., Bellanger, A.P., Roussel, S., Grenouillet, F., Sornin, S., Piarroux, R., Dalphin,
507 J.C., Millon, L., 2009. Indoor mold concentration in Eastern France. *Indoor Air* 19,
508 446–453. <https://doi.org/10.1111/j.1600-0668.2009.00598.x>

509 Scherer, E., Rocchi, S., Reboux, G., Vandentorren, S., Roussel, S., Vacheyrou, M., Raheison,
510 C., Millon, L., 2014. qPCR standard operating procedure for measuring
511 microorganisms in dust from dwellings in large cohort studies. *Sci. Total Environ.*
512 466–467, 716–724. <https://doi.org/10.1016/j.scitotenv.2013.07.054>

513 Shendell, D.G., Mizan, S.S., Yamamoto, N., Peccia, J., 2012. Associations between
514 quantitative measures of fungi in home floor dust and lung function among older
515 adults with chronic respiratory disease: a pilot study. *J. Asthma* 49, 502–509.
516 <https://doi.org/10.3109/02770903.2012.682633>

517 Soler-Cataluña, J.J., Martínez-García, M.Á., Sánchez, P.R., Salcedo, E., Navarro, M.,
518 Ochando, R., 2005. Severe acute exacerbations and mortality in patients with chronic
519 obstructive pulmonary disease. *Thorax* 60, 925–931.
520 <https://doi.org/10.1136/thx.2005.040527>

521 Tiew, P.Y., Ko, F.W.S., Pang, S.L., Matta, S.A., Sio, Y.Y., Poh, M.E., Lau, K.J.X., Mac
522 Aogáin, M., Jaggi, T.K., Ivan, F.X., Gaultier, N.E., Uchida, A., Drautz-Moses, D.I.,
523 Xu, H., Koh, M.S., Hui, D.S.C., Tee, A., Abisheganaden, J.A., Schuster, S.C., Chew,
524 F.T., Chotirmall, S.H., 2020. Environmental fungal sensitisation associates with
525 poorer clinical outcomes in COPD. *Eur. Respir. J.*
526 <https://doi.org/10.1183/13993003.00418-2020>

527 Uffredi, M.L., Mangiapan, G., Cadranel, J., Kac, G., 2003. Significance of *Aspergillus*
528 *fumigatus* isolation from respiratory specimens of nongranulocytopenic patients. *Eur.*
529 *J. Clin. Microbiol. Infect. Dis.* 22, 457–462. <https://doi.org/10.1007/s10096-003-0970->
530 y

531 Vesper, S., McKinstry, C., Ashley, P., Haugland, R., Yeatts, K., Bradham, K., Svendsen, E.,
532 2007. Quantitative PCR analysis of molds in the dust from homes of asthmatic
533 children in North Carolina. *J. Environ. Monit. JEM* 9, 826–830.
534 <https://doi.org/10.1039/b704359g>

535 Viegas, C., Faria, T., Caetano, L.A., Carolino, E., Quintal-Gomes, A., Twarużek, M., Kosicki,
536 R., Viegas, S., 2019. Characterization of Occupational Exposure To Fungal Burden in
537 Portuguese Bakeries. *Microorganisms* 7.
538 <https://doi.org/10.3390/microorganisms7080234>

539 Williams, B., Douglas, P., Roca Barcelo, A., Hansell, A.L., Hayes, E., 2019. Estimating
540 *Aspergillus fumigatus* exposure from outdoor composting activities in England
541 between 2005 and 14. *Waste Manag.* 84, 235–244.
542 <https://doi.org/10.1016/j.wasman.2018.11.044>

543

544

545 **Figure legends**

546 **Fig. 1.** Flow diagram of patient recruitment and follow-up

547

548 **Fig. 2.** Kinetics of anti-*Aspergillus* antibodies in patients with aspergillosis diagnosed at the
549 final visit

550

551 **Fig. 3.** Association between FEV1 and *Aspergillus* biomarkers

552 FEV1 was compared between patients with positive or negative *Aspergillus* section *Fumigati*
553 culture in sputum (A) and between patients with or without anti-*Aspergillus* antibodies (B) at
554 inclusion.

555

556 **Fig. 4.** Association between domestic mould contamination levels (total or *Penicillium* spp.
557 counts) and *Aspergillus* biomarkers

558 Total mould (A-C) and *Penicillium* spp. (D-F) concentrations in EDCs were compared
559 between patients with positive or negative *Aspergillus* section *Fumigati* culture in sputum at
560 inclusion and between patients with or without anti-*Aspergillus* antibodies at inclusion, and at
561 final visit.

562

563 **Fig. 5.** Association between domestic mould contamination levels (total or *Aspergillus* section
564 *Fumigati* counts) and COPD stage (A) or number of exacerbations (B)

565

Fig. 1





Fig. 2

N°10
Anti-*A. fumigatus* precipitins



N°52
Anti-*A. fumigatus* precipitins




Fig. 3





Fig. 4




Fig. 5




Table 1

Predictors of *Aspergillus* section *Fumigati* positive sputum culture or anti-*Aspergillus* antibody detection during exacerbation

	<i>Aspergillus</i> section <i>Fumigati</i> detection in sputum			Detection of anti- <i>Aspergillus</i> antibodies		
	Positive	Negative	P-values	Positive	Negative	P-values
Patients (n)	10	49		20	42	
Men (%)	80.0	69.4	0.40	75.0	71.4	0.51
Age	61.1±2.4	63.9±1.7	0.23	61.5±1.7	64.6±1.9	0.14
Body mass index (BMI) (kg/m ²)	28.7±1.6	24.4±0.9	0.01	23.0±1.1	26.2±1.1	0.03
Pack years	67.9±12.1	61.5±4.7	0.29	71.8±6.7	59.3±5.3	0.08
Current smokers (%)	30.0	26.5	0.55	20.0	30.9	0.28
≥ 1 exacerbation with hospitalization/year in the 2 previous years (%)	66.7	66.7	0.64	64.3	70	0.63
GOLD II/III/IV	0/50.0/50.0	18.4/44.9/36.7	0.32	5.0/35.0/60.0	19.0/50.0/31.0	0.07
Inhaled corticosteroid (%)	70	81.6	0.33	95.0	71.4	0.03
FEV1 (mean %)	28.7 ± 3.3	37.3 ± 2.4	0.06	28.5 ± 3.3	39.1 ± 2.4	0.01
Confirmed bacterial exacerbation (%)	40.0	32.7	0.52	45.0	28.6	0.16

Data are presented as n (%), mean±SEM, unless otherwise stated. Bold type represents statistical significance at p<0.05.

Table 2

Mould species frequency and number of colonies in electrostatic dust collectors

Species	Frequency (%)	Median CFUs/cm ² (min-max)
<i>Penicillium</i> spp.	90.2	3.1 (0.6-64.2)
<i>Cladosporium</i> spp.	64.7	1.9 (0.6-15.4)
<i>Aspergillus</i> section <i>Fumigati</i>	49.0	1.2 (0.6-11.7)
<i>Aspergillus</i> series <i>Versicolores</i>	39.2	0.9 (0.6-24.7)
<i>Mucorales</i>	33.3	1.2 (0.6-9.3)
- <i>Mucor</i> spp.	15.7	0.9 (0.6-1.9)
- <i>Rhizopus</i> spp.	13.7	0.6 (0.6-1.2)
- <i>Lichtheimia</i> spp.	2.0	0.6 (0.6-0.6)
- <i>Cunninghamella</i> spp.	2.0	0.6 (0.6-0.6)
- Other mucorales	7.8	0.9 (0.6-9.3)
<i>Aspergillus</i> section <i>Nigri</i>	25.5	1.2 (0.6-6.2)
<i>Aspergillus</i> spp.	25.5	1.2 (0.6-54.9)
<i>Alternaria</i> spp.	25.5	1.2 (0.6-4.3)
<i>Fusarium</i> spp.	19.6	1.5 (0.6-34.6)
<i>Scopulariopsis</i> spp.	17.6	0.6 (0.6-1.9)
<i>Wallemia</i> spp.	11.8	0.9 (0.6-3.7)
<i>Geotrichum</i> spp.	11.8	0.6 (0.6-1.2)
<i>Aspergillus</i> section <i>Flavi</i>	9.8	3.1 (0.6-7.4)
<i>Paecilomyces</i> spp.	9.8	0.6 (0.6-0.6)
<i>Trichoderma</i> spp.	9.8	0.6 (0.6-0.6)
<i>Botrytis</i> spp.	5.9	0.6 (0.6-1.2)
<i>Ulocladium</i> spp.	5.9	0.6 (0.6-0.6)
<i>Aspergillus</i> series <i>Nidulantes</i>	3.9	0.9 (0.6-1.2)
Other fungi	≤2.0	NA

Table 3

Frequency of *Aspergillus* section *Fumigati* detection in sputum and anti-*Aspergillus* antibodies in patients with total mould contamination level of \leq or $>$ 30 CFUs/cm² at inclusion and final visit

	Total counts		
	\leq 30 CFUs/cm ²	$>$ 30 CFUs/cm ²	P-values
<i>Aspergillus</i> section <i>Fumigati</i> detection in sputum			
Inclusion (n=48)	7/41 (17%)	2/7 (29%)	0.39
Final visit (n=25)	0/20 (0%)	1/5 (20%)	0.20
Detection of anti- <i>Aspergillus</i> antibodies			
Inclusion (n=51)	13/43 (30%)	5/8 (63%)	0.09
Final visit (n=32)	6/25 (24%)	5/7 (71%)	0.03

Bold type represents statistical significance at $p < 0.05$.