



HAL
open science

Manipulable outcomes within the class of scoring voting rules

Mostapha Diss, Boris Tselikhovskiy

► **To cite this version:**

Mostapha Diss, Boris Tselikhovskiy. Manipulable outcomes within the class of scoring voting rules. 2024. hal-04543626

HAL Id: hal-04543626

<https://univ-fcomte.hal.science/hal-04543626>

Preprint submitted on 12 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

crese

CENTRE DE RECHERCHE
SUR LES STRATÉGIES ÉCONOMIQUES

Manipulable outcomes within the class of scoring voting rules

MOSTAPHA DISS, BORIS TSVELIKHOVSKIY

November 2020

Working paper No. 2020 – 08

CRESE

30, avenue de l'Observatoire
25009 Besançon
France
<http://crese.univ-fcomte.fr/>

The views expressed are those of the authors
and do not necessarily reflect those of CRESE.

UFR SJE PG 

Sciences juridiques économiques
politiques et de gestion

**UNIVERSITÉ DE
FRANCHE-COMTÉ**

Manipulable outcomes within the class of scoring voting rules*

Mostapha Diss[†] Boris Tsvetikhovskiy[‡]

November 9, 2020

Abstract

Coalitional manipulation in voting is considered to be any scenario in which a group of voters decide to misrepresent their vote in order to secure an outcome they all prefer to the first outcome of the election when they vote honestly. The present paper is devoted to study coalitional manipulability within the class of scoring voting rules. For any such rule and any number of alternatives, we introduce a new approach allowing to characterize all the outcomes that can be manipulable by a coalition of voters. This gives us the possibility to find the probability of manipulable outcomes for some well-studied scoring voting rules in the case of small number of alternatives and large electorates under a well-known assumption on individual preference profiles.

Keywords: Voting; Scoring Rules; Coalition; Strategic Manipulation; Probability.

JEL classification: D71, D72

1 Introduction

Since the seminal papers of [Gibbard \(1973\)](#) and [Satterthwaite \(1975\)](#) who proved that every non-dictatorial social choice rule can be manipulated in the presence of at least three alterna-

*The authors would like to thank Eric Kamwa for comments and helpful discussions. The first author gratefully acknowledges the financial supports from Université de Lyon (project INDEPTH Scientific Breakthrough Program of IDEX Lyon) within the program Investissement d'Avenir (ANR-16-IDEX-0005) and from Université de Franche-Comté within the program Chrysalide-2020. The second author wishes to express his gratitude to F. Aleskerov for introducing him to the subject and constant guidance on the early stages of development of this project. He also acknowledges the support of the International Laboratory of Decision Choice and Analysis (National Research University Higher School of Economics).

[†]CRESE EA3190, Univ. Bourgogne Franche-Comté, F-25000 Besançon, France. Email: mostapha.diss@univ-fcomte.fr.

[‡]Department of Mathematics, Northeastern University, Boston, MA, 02115, USA. Email: tsvelikhovskiy.b@northeastern.edu.

tives, the problem of coalitional manipulation has received a lot of attention in recent decades in social choice theory. Broadly speaking, a given social choice rule is called *coalitionally manipulable* if there exists a given list of voting preferences and a coalition of voters,¹ such that the preferences of all the voters outside the coalition remain the same, while the preferences of voters within the coalition can be altered in such a way that the winner changes and each of the voters from the coalition is ‘happy about the change’.

Scoring voting rules, also called *positional voting rules*, have attracted a considerable amount of attention in the literature dealing with manipulation. This class of voting rules can be defined as follows: each voter’s preference must be a vector that gives a number of points that the voter assigns to each alternative according to his or her position in the voter’s preference. The points assigned by all voters are summed, and the winning alternative has the highest number of points. A number of studies has been conducted on the evaluation of the degree of manipulability of various social choice rules, i.e., the extent to which social choice rules are manipulable by a coalition of voters or by an individual voter. The reader may refer, for instance, to [Aleskerov and Kurbanov \(1999\)](#), [Chamberlin \(1985\)](#), [Diss \(2015\)](#), [El Ouafdi et al. \(2020a\)](#), [El Ouafdi et al. \(2020b\)](#), [Favardin and Lepelley \(2006\)](#), [Favardin et al. \(2002\)](#), [Gehrlein et al. \(2013\)](#), [Kamwa and Moyouwou \(2020\)](#), [Kelly \(1993\)](#), [Kim and Roush \(1996\)](#), [Lepelley and Mbih \(1994\)](#), [Lepelley et al. \(2008\)](#), [Moyouwou and Tchantcho \(2017\)](#), [Nitzan \(1985\)](#), [Peleg \(1979\)](#), [Pritchard and Wilson \(2007a\)](#), [Pritchard and Wilson \(2007b\)](#), [Saari \(1990\)](#), and [Schürmann \(2013\)](#). The methodology used in this literature consists first in characterizing the specific conditions that must be required for a given voting rule to be manipulable by a coalition of voters. The final step consists in the evaluation of the (theoretical) probability of this situation under various assumptions on voters’ preferences. For more details on those probabilistic assumptions and their use in social choice theory, the reader can refer to [Diss and Merlin \(2020\)](#), [Gehrlein and Lepelley \(2017\)](#), [Gehrlein and Lepelley \(2011\)](#).

However, as pointed out in a recent paper by [El Ouafdi et al. \(2020a\)](#), with some notable exceptions the results appearing in the literature only deal with three-alternative elections, not because it is the most interesting case, but due to the difficulties arising when considering more than three alternatives. The main goal of this paper is to provide a significant improvement in this direction. This is achieved via presenting systems of linear inequalities, which determine manipulable profiles, that is the lists defining the vote of all voters taking part in the decision process, for the whole class of scoring voting rules independently on the number of alternatives. More precisely, we focus on a detailed exposition of the new approach for obtaining the list of linear inequalities a profile satisfies if and only if it is manipulable in the case of $m \geq 3$ alternatives for all scoring rules.

The paper is organized as follows. Section 2 describes the basic framework. The core of the paper is Section 3, where the main results are presented. We start off with providing a simplified illustration of how our methodology works in the case of three-alternative elections. Then, the precise list of inequalities for the whole class of scoring rules is pre-

¹A coalition is defined as any non-empty subset of voters.

sented. Attention will be focused on the limiting case where the number of voters tends to infinity. This will enable us to complete the existing literature dealing with the probability of manipulable outcomes by providing in Section 4 the corresponding values for the three most studied scoring rules in this literature which are Plurality, Antiplurality and Borda rules in the presence of $m = 4$ and $m = 5$ alternatives under the well-known Impartial and Anonymous Culture assumption (defined later). The results are approximate since they are obtained by Monte-Carlo simulations, but with a high degree of precision, since we used profiles of cardinality $8 \cdot 10^6$ for $m = 4$ and $8 \cdot 10^5$ for $m = 5$ alternatives. To the best of our knowledge none of those results have appeared in the literature, with the exception of the Plurality rule and $m = 4$ (see El Ouafdi et al. (2020a)). The new results clearly indicate that among the three positional rules under consideration Antiplurality has a much lesser degree of manipulability (see Tables 5, 6, and 7). The last section presents our conclusions.

2 Preliminary definitions

Let us assume that the number of voters/individuals is denoted by n and the number of alternatives/candidates by m . The voters' preferences are assumed to be linear orders which means that voters rank alternatives from most preferred to least preferred and indifference is not allowed. An *anonymous preference profile*, hereafter simply called a *profile*, is an $m!$ -tuple of non-negative integer numbers $(n_1, n_2, \dots, n_{m!})$ such that $n_i \geq 0$ for all $m! \geq i \geq 1$ and $\sum_{i=1}^{m!} n_i = n$.² Each n_i is equal to the number of voters with preferences of type i . In the limit as $n \rightarrow \infty$, we consider the normalized profile vectors of the form $p = (p_1, \dots, p_{m!})$ with each $p_i = \frac{n_i}{n} \geq 0$ and $\sum_{i=1}^{m!} p_i = 1$. Throughout the paper, the alternatives will be called A_1, A_2, \dots, A_m and the types of possible preference rankings of the alternatives will always be listed following an increasing order of j in the notation A_j . In addition, the proportion of individuals having each possible preference ranking will be denoted accordingly. For instance, in the case of $m = 3$ alternatives, the possible preference rankings are denoted by $(A_1, A_2, A_3) p_1$, $(A_1, A_3, A_2) p_2$, $(A_2, A_1, A_3) p_3$, $(A_2, A_3, A_1) p_4$, $(A_3, A_1, A_2) p_5$ and $(A_3, A_2, A_1) p_6$. The notation $(A_1, A_2, A_3) p_1$, for instance, means that a fraction p_1 of individuals have preferences with A_1 being most preferred, A_3 being least preferred and with A_2 being ranked between them.

In the case of m alternatives and individual preferences expressed as linear orders, we can define the class of scoring voting rules as follows:

Definition 2.1. In the case of linear orders on m alternatives, every scoring voting rule can be defined by the *weight vector* $w = (w_1, w_2, \dots, w_m)$ such that $w_1 \geq w_2 \geq \dots \geq w_m$ and $w_1 > w_m$. It means that every time an alternative is ranked r -th by one voter it obtains w_r points; afterwards we select the alternative(s) obtaining the greatest aggregated score.

²This is also known as a *voting situation* in the literature.

Given this definition, we are now ready to describe three well-known scoring voting rules. Under Plurality rule each voter has one vote which he/she can cast for any one of the m alternatives, i.e., $w = (1, 0, \dots, 0)$. Under Borda rule, we select the alternative with the highest Borda score such that each first-place vote is worth $m - 1$ points, each second-place vote is worth $m - 2$ points, and so on until 0 points are assigned to each last-place vote, i.e., $w = (m - 1, m - 2, \dots, 1, 0)$. Under Antiplurality rule, we return the alternative with the highest aggregated score such that each voter assigns one point for any one of the $m - 1$ best ranked alternatives, i.e., $w = (1, 1, \dots, 1, 0)$.

Note that in the presence of three alternatives $A_1, A_2,$ and $A_3,$ we see that any general scoring rule $w = (w_1, w_2, w_3)$ is equivalent to the normalized scoring vector $(1, \lambda, 0)$ with $0 \leq \lambda \leq 1$. This is obtained by subtracting w_3 and dividing by $w_1 - w_3$ any component of w . The well-known scoring rules become the Plurality rule with $\lambda = 0$, the Borda rule with $\lambda = 0.5$, and the Antiplurality rule with $\lambda = 1$.

As already mentioned, we focus in this paper on the coalitional manipulation which can be defined as follows:

Definition 2.2. The outcome of an election is said to be *coalitionally manipulable* if there exists an alternative such that all members of the electorate for whom she is preferable over the winning alternative can change their preferences in such a way that this alternative wins the election. The preferences of the rest of the electorate remain the same.

We mainly focus in this paper on the manipulation by coalitions of maximal sizes but we will also show that the approach that we consider can be adapted when manipulation by coalitions of smaller sizes of the electorate is considered.

We say that the initial arrangement is (A_1, A_2, \dots, A_m) if the collective ranking of the alternatives before manipulation is: A_1 first, A_2 second and so on until alternative A_m which is ranked last. Let us consider an example in order to highlight the notion of manipulation.

Example 2.3. Let $m = 3$ and the scoring rule be Borda rule, i.e., $w = (2, 1, 0)$. To show a manipulable outcome, we set $p_1 = 5/9, p_3 = 4/9,$ and $p_j = 0$ for $j \neq 1, 3$. By the definition of Borda rule, the initial arrangement is (A_1, A_2, A_3) since alternative A_1 wins and gets $14/9$ points, A_2 gets $13/9$ points, and finally A_3 gets 0 points. But, if voters having the preference (A_2, A_3, A_1) chose to vote (A_2, A_1, A_3) instead, then the final result would be in favor of alternative A_2 since A_1 gets $10/9$ points, A_2 gets $13/9$ points, and A_3 gets $4/9$ points.

With a given initial arrangement, $m - 1$ different coalitions can be formed. This is due to the fact that for each alternative, who has not won the election, there may be a group of voters who rank it higher than the winner. As any coalition is determined by the alternative it would prefer to have as the winner of the election, the alternatives' names will be used for the notations of the coalitions hereafter. Alternative A_i will be called *unifying* for the coalition denoted by Coal_{A_i} that would prefer to have A_i as the winner and let us denote by \mathfrak{C}_{A_i} the proportion of individuals in this coalition. In the case of $m = 3$ alternatives, for instance,

two coalitions can be formed since the unifying alternative can be any alternative except the winner. In this case, when the winner is alternative A_1 , we have that alternative A_2 is unifying for the coalition Coal_{A_2} formed by voters with preferences (A_2, A_1, A_3) , (A_2, A_3, A_1) and (A_3, A_2, A_1) and alternative A_3 is unifying for the coalition Coal_{A_3} formed by voters with preferences (A_3, A_1, A_2) , (A_3, A_2, A_1) and (A_2, A_3, A_1) . It is clear that the preferences (A_2, A_3, A_1) and (A_3, A_2, A_1) participate in both coalitions Coal_{A_2} and Coal_{A_3} .

We introduce now the concept of *intermediate preferences*.

Definition 2.4. *Intermediate preferences* are preferences of the form $(*, *, \dots, *)$ with each "*" symbol being either an alternative's name or the "?" mark, the latter represents the fact that its corresponding rank has yet to be assigned by the voter.

Typical examples of intermediate preferences in three-alternative elections are $(?, ?, ?)$ where the concerned voter is undecided on all of the three positions and $(A_1, ?, ?)$ where the concerned voter ranks alternative A_1 at the first position and he/she is undecided on the second and third positions in his/her preference.

Let us now introduce $d(A_k, A_i)$, with $i \neq k$, which denotes the difference in points between the unifying alternative A_k and alternative A_i prior to the coalition participants' arrangement of all the places except the first. In the case of three alternatives, when the winner is alternative A_1 for instance, individuals of Coal_{A_2} (of proportion $\mathfrak{C}_{A_2} = p_3 + p_4 + p_6$) vote for alternative A_2 , i.e., the unifying alternative of that coalition. In addition, those individuals only assign weight 1 to their first preferred alternative and zero weight to the other alternatives prior to the coalition participants' arrangement of the second and third places. In other words, those individuals have an intermediate preference of type $(A_2, ?, ?)$. Note that the other individuals (of proportion $p_1 + p_2 + p_5$) assign all their points and vote sincerely when we focus on the manipulation in favor of A_2 . As a consequence, the score of alternative A_2 is $\lambda p_1 + p_3 + p_4 + p_6$ and the score of alternative A_1 is $p_1 + p_2 + \lambda p_5$. This leads to $d(A_2, A_1) = p_3 + p_4 + p_6 - p_2 + (\lambda - 1)p_1 - p_5$.

3 Conditions describing manipulability

In this section we show that for every coalition there exists a system of inequalities which characterize the set of all the profiles that are manipulable by it. In order to make a simplified illustration of the general case that we will provide in Theorem 3.1, let us start first by the case of three-alternative elections. Let again the alternatives be referred to as A_1 , A_2 , and A_3 and consider the representation $w = (1, \lambda, 0)$, with $0 \leq \lambda \leq 1$, that defines all possible scoring rules with three alternatives. Using all the above definitions, we are ready to provide the list of inequalities which characterizes the set of all the profiles $p = (p_1, \dots, p_6)$ that are manipulable by a coalition of voters for three alternatives. The first system indicates the conditions under which the manipulation is in favour of A_2 ; the second system concerns

manipulation in favour of A_3 .

$$\begin{cases}
 p_i \geq 0, i \in \{1, 2, \dots, 6\} \\
 p_1 + p_2 + p_3 + p_4 + p_5 + p_6 = 1 \\
 (1 - \lambda)p_1 + p_2 + (\lambda - 1)p_3 - p_4 + \lambda p_5 - \lambda p_6 > 0, & A_1 \text{ beats } A_2 \\
 \lambda p_1 - \lambda p_2 + p_3 + (1 - \lambda)p_4 - p_5 + (\lambda - 1)p_6 > 0, & A_2 \text{ beats } A_3 \\
 p_3 + p_4 + p_6 - p_2 + (\lambda - 1)p_1 - \lambda p_5 > 0, & d(A_2, A_1) > 0 \\
 (2 - \lambda)(p_3 + p_4 + p_6) + (2\lambda - 1)p_1 - (1 + \lambda)p_2 - (1 + \lambda)p_5 > 0, & d(A_2, A_1) + d(A_2, A_3) > \lambda \mathfrak{C}_{A_2}
 \end{cases}$$

$$\begin{cases}
 p_i \geq 0, i \in \{1, 2, \dots, 6\} \\
 p_1 + p_2 + p_3 + p_4 + p_5 + p_6 = 1 \\
 (1 - \lambda)p_1 + p_2 + (\lambda - 1)p_3 - p_4 + \lambda p_5 - \lambda p_6 > 0, & A_1 \text{ beats } A_2 \\
 \lambda p_1 - \lambda p_2 + p_3 + (1 - \lambda)p_4 - p_5 + (\lambda - 1)p_6 > 0, & A_2 \text{ beats } A_3 \\
 p_4 + p_5 + p_6 - p_1 + (\lambda - 1)p_2 - \lambda p_3 > 0, & d(A_3, A_1) > 0 \\
 p_4 + p_5 + p_6 - p_3 + \lambda p_2 - \lambda p_1 > 0, & d(A_3, A_2) > 0 \\
 (2 - \lambda)(p_4 + p_5 + p_6) - (1 + \lambda)p_1 + (2\lambda - 1)p_2 - (1 + \lambda)p_3 > 0, & d(A_3, A_1) + d(A_3, A_2) > \lambda \mathfrak{C}_{A_3}.
 \end{cases}
 \tag{3.1}$$

Without loss of generality we assume that the initial arrangement is (A_1, A_2, A_3) . This means that A_1 has more total points than A_2 who in turn has more total points than A_3 . This explains the two first inequalities of each system. Recall that when the winner is alternative A_1 we have that alternative A_2 is unifying for the coalition Coal_{A_2} formed by voters with preferences $(A_2, A_1, A_3), (A_2, A_3, A_1)$, and (A_3, A_2, A_1) . If we count the number of points for each alternative prior to the coalition participants' arrangement of all the places except the first which is given to the unifying alternative A_k (i.e., having intermediate preferences of the form $(A_k, ?, ?)$), the unifying alternative A_k must have the largest number of points. Thus, the two inequalities $d(A_k, A_i) > 0$, $i \neq k$ are necessary with $k = 2$ and $k = 3$ for the first and the second systems, respectively. Note that the inequality called ' A_2 beats A_3 ' in the first system leads to $d(A_2, A_3) > 0$. The next step is to understand when individuals of the coalition unified by an alternative can successfully manipulate the election. Actually, all the freedom they possess at this point is to give λ points from each participant of the coalition (by assigning the second place) to one of the remaining two alternatives without violating the condition that their alternative has more points. This can be done if and only if the sum of the differences in points between the unifying alternative and each of the remaining two, prior to the coalition members' making a choice of their second most preferable alternative, is greater than λ times the number of people in the coalition. This is equivalent to $d(A_2, A_1) + d(A_2, A_3) > \lambda \mathfrak{C}_{A_2}$ and $d(A_3, A_1) + d(A_3, A_2) > \lambda \mathfrak{C}_{A_3}$ in the first and the second systems, respectively. Recalling that we consider a manipulation by coalitions of maximal sizes (i.e., $\mathfrak{C}_{A_2} = p_3 + p_4 + p_6$ and $\mathfrak{C}_{A_3} = p_4 + p_5 + p_6$), this leads to the last inequality of every system. That analysis gives first insights on the validity of our approach since the systems described above are exactly the same as the ones already obtained by [Moyouwou and Tchantcho \(2017\)](#) who determined necessary and sufficient conditions

for a given profile to be coalitionally manipulable under a general scoring rule with three alternatives.

We consider now the general case of m alternatives. Recall that the unifying alternative can be any alternative except the winner so that $m - 1$ coalitions can be formed. Here again we consider the coalitions of maximal sizes, i.e., each coalition consists of all voters with the unifying alternative ranked higher than the winner in their preferences. The set of such preferences has cardinality $\frac{m!}{2}$.

Theorem 3.1. *Let $m \geq 3$ and suppose that the initial arrangement is (A_1, A_2, \dots, A_m) under the positional voting rule having weight vector $w = (w_1, w_2, \dots, w_m)$. A profile $p = (p_1, \dots, p_{m!})$ is manipulable by the coalition with unifying alternative A_k if and only if p satisfies the following system of inequalities:*

$$\left\{ \begin{array}{l} \text{Preliminary Inequalities (PI)} \\ \left\{ \begin{array}{l} \sum_{i=1}^{m!} p_i = 1 \\ p_i \geq 0 \\ \text{The initial arrangement is } (A_1, A_2, \dots, A_m) \end{array} \right. \\ \\ \text{Strategic Inequalities (SI)} \\ \left\{ \begin{array}{l} \sum_{i \neq k} d(A_k, A_i) > (w_2 + \dots + w_{m-1} + w_m) \mathfrak{C}_{A_k} \\ \sum_{i \neq k} d(A_k, A_i) - M_1 > (w_3 + \dots + w_{m-1} + w_m) \mathfrak{C}_{A_k} \\ \dots \\ \sum_{i \neq k} d(A_k, A_i) - \sum_{j=1}^{m-4} M_j > (w_{m-2} + w_{m-1} + w_m) \mathfrak{C}_{A_k} \\ \sum_{i \neq k} d(A_k, A_i) - \sum_{j=1}^{m-3} M_j > (w_{m-1} + w_m) \mathfrak{C}_{A_k} \\ M_{m-1} > w_m \mathfrak{C}_{A_k}, \end{array} \right. \end{array} \right. \quad (3.2)$$

where $M_1 = \max\{d(A_k, A_i) \mid i \neq k\}$, $M_2 = \max_2\{d(A_k, A_i) \mid i \neq k\}$, \dots , $M_{m-1} = \max_{m-1}\{d(A_k, A_i) \mid i \neq k\}$. The notation $\max_s\{d(A_k, A_i) \mid i \neq k\}$ stands for the s^{th} largest value in $\{d(A_k, A_i) \mid i \neq k\}$.

Proof. First let us show that the SI inequalities in (3.2) are necessary. This will be checked by induction on m , the base being $m = 3$. Notice that it is clearly reasonable for all the members of the coalition to start filling their profiles by updating their intermediate preferences to $(A_k, ?, \dots, ?)$. The first inequality in the SI system simply guarantees that after that update the coalition participants can distribute the remaining votes they have to give without violating the fact that A_k has more points than any other alternatives with no restrictions on the distribution of the votes. The second to last inequalities of the SI system can be seen to be those for the positional rule with $w' = (w_1, w_3, \dots, w_{m-1}, w_m)$ and $m - 1$ alternatives

(alternative A_s with $M_1 = d(A_k, A_s)$ is eliminated). The statement follows.

Now we establish the sufficiency of inequalities (3.2). We argue by induction on the number of alternatives m , the base being $m = 3$. The sufficiency of inequalities (3.2) is established via the step by step procedure of updating the intermediate preferences of coalition participants presented below.

Step 1. All the members of the coalition start filling their preferences by putting A_k as the most preferable alternative. At this point the preferences of all participants of Coal_{A_k} are identical and equal to $(A_k, ?, \dots, ?)$, where the question marks stand for the places yet to be assigned.

Step 2. Let A_s be the alternative with $d(A_k, A_s) = M_1$. We will distinguish between two cases:

case (a) $M_1 > w_2 \mathfrak{C}_{A_k}$. Then all participants of Coal_{A_k} update their intermediate preferences to $(A_k, A_s, ?, \dots, ?)$ and the list of inequalities in the SI system of (3.2) without the second one from the top is easily seen to be the one for the positional rule with $m - 1$ alternatives and weight vector (w_1, w_3, \dots, w_m) ;

case (b) $w_\sigma \mathfrak{C}_{A_k} < M_1 < w_{\sigma-1} \mathfrak{C}_{A_k}$ for some $2 < \sigma \leq m$. We would like to point out that the case $M_1 < w_m \mathfrak{C}_{A_k}$ can never occur as it would imply $\sum_{i \neq k} d(A_k, A_i) < (m-1)w_m \mathfrak{C}_{A_k} < (w_2 + \dots + w_{m-1} + w_m) \mathfrak{C}_{A_k}$ and violate the first (SI) inequality.

Let $\varepsilon > 0$ be an arbitrary small positive constant, satisfying

(\star) ε does not exceed the smallest difference between the left-hand side and right-hand side of the SI inequalities of (3.2).

It will serve as the remaining difference in points between A_k and A_s after the update of preferences described below. Set $t := \frac{M_1 - w_\sigma \mathfrak{C}_{A_k}}{w_{\sigma-1} - w_\sigma} - \varepsilon$ and let t ‘participants’ update their intermediate preferences to $(A_k, ?, \dots, ?, A_s, ?, \dots, ?)$ and the remaining $\mathfrak{C}_{A_k} - t$ to $(A_k, ?, \dots, ?, A_s, ?, \dots, ?)$. Introduce the weight

$$w_{\sigma-1, \sigma} := \frac{(\mathfrak{C}_{A_k} - t)w_{\sigma-1} + tw_\sigma}{\mathfrak{C}_{A_k}} \quad (3.3)$$

and set

$$w' = (w_1, \dots, w_{\sigma-2}, w_{\sigma-1, \sigma}, w_{\sigma+1}, \dots, w_m). \quad (3.4)$$

After updates of coalition participants’ intermediate preferences and elimination of alternative A_s , the system of SI inequalities in (3.2) becomes the one for the

positional rule with $m - 1$ alternatives and weight vector w' :

$$\left\{ \begin{array}{l} \sum_{i \neq k, s} d(A_k, A_i) > (w_2 + \dots + w_{\sigma-1, \sigma} + \dots + w_m) \mathfrak{C}_{A_k} \\ \sum_{i \neq k, s} d(A_k, A_i) - M_2 > (w_3 + \dots + w_{\sigma-1, \sigma} + \dots + w_m) \mathfrak{C}_{A_k} \\ \dots \\ \sum_{i \neq k, s} d(A_k, A_i) - \sum_{j=2}^{\sigma-2} M_j > (w_{\sigma-1, \sigma} + \dots + w_m) \mathfrak{C}_{A_k} \\ \sum_{i \neq k, s} d(A_k, A_i) - \sum_{j=2}^{\sigma-1} M_j > (w_{\sigma+1} + \dots + w_m) \mathfrak{C}_{A_k} \\ \dots \\ \sum_{i \neq k, s} d(A_k, A_i) - \sum_{j=2}^{m-4} M_j > (w_{m-2} + w_{m-1} + w_m) \mathfrak{C}_{A_k} \\ \sum_{i \neq k, s} d(A_k, A_i) - \sum_{j=2}^{m-3} M_j > (w_{m-1} + w_m) \mathfrak{C}_{A_k} \\ M_{m-1} > w_m \mathfrak{C}_{A_k}, \end{array} \right. \quad (3.5)$$

which, using that $M_1 = d(A_k, A_s) = (w_{\sigma-1} + w_{\sigma} - w_{\sigma-1, \sigma}) \mathfrak{C}_{A_k}$, can be rewritten as

$$\left\{ \begin{array}{l} \sum_{i \neq k} d(A_k, A_i) > (w_2 + \dots + w_{\sigma-1} + w_{\sigma} + \dots + w_m) \mathfrak{C}_{A_k} + \varepsilon \\ \sum_{i \neq k} d(A_k, A_i) - M_2 > (w_3 + \dots + w_{\sigma-1} + w_{\sigma} + \dots + w_m) \mathfrak{C}_{A_k} + \varepsilon \\ \dots \\ \sum_{i \neq k} d(A_k, A_i) - \sum_{j=2}^{\sigma-2} M_j > (w_{\sigma-1} + w_{\sigma} + \dots + w_m) \mathfrak{C}_{A_k} + \varepsilon \\ \sum_{i \neq k} d(A_k, A_i) - \sum_{j=2}^{\sigma-1} M_j > (w_{\sigma+1} + \dots + w_m) \mathfrak{C}_{A_k} \\ \dots \\ \sum_{i \neq k} d(A_k, A_i) - \sum_{j=1}^{m-4} M_j > (w_{m-2} + w_{m-1} + w_m) \mathfrak{C}_{A_k} \\ \sum_{i \neq k} d(A_k, A_i) - \sum_{j=1}^{m-3} M_j > (w_{m-1} + w_m) \mathfrak{C}_{A_k} \\ M_{m-1} > w_m \mathfrak{C}_{A_k} \end{array} \right. \quad (3.6)$$

Recalling that M_1 is greater than any other M_i by definition and using the (\star) assumption on ε , we see that each of the inequalities in (3.6) follows from the corresponding SI inequality in (3.2).

□

Let us now illustrate the results of this theorem using an example.

Example 3.2. Consider the Borda rule with weight vector $w = (3, 2, 1, 0)$ for $m = 4$ alternatives A_1, A_2, A_3 , and A_4 and the voting profile given in Table 1 below. In this example, the initial arrangement is (A_1, A_2, A_3, A_4) as it is shown in the last row. We would like to determine if the chosen profile is vulnerable to manipulability by the largest possible coalition with unifying alternative A_2 , i.e., Coal_{A_2} of cardinality $\mathfrak{C}_{A_2} = p_7 + p_8 + p_{15} = 5/9$.

Table 1: Voting profile and corresponding point distribution

Preference	Fraction	A_1	A_2	A_3	A_4
(A_2, A_1, A_3, A_4)	$p_7 = 2/9$	$4/9$	$6/9$	$2/9$	0
(A_2, A_1, A_4, A_3)	$p_8 = 2/9$	$4/9$	$6/9$	0	$2/9$
(A_3, A_1, A_4, A_2)	$p_{14} = 1/9$	$2/9$	0	$3/9$	$1/9$
(A_3, A_2, A_1, A_4)	$p_{15} = 1/9$	$1/9$	$2/9$	$3/9$	0
(A_3, A_4, A_1, A_2)	$p_{17} = 1/9$	$1/9$	0	$3/9$	$2/9$
(A_4, A_1, A_3, A_2)	$p_{20} = 2/9$	$4/9$	0	$2/9$	$6/9$
Other	$p_i = 0$	0	0	0	0
\sum	1	$16/9$	$14/9$	$13/9$	$11/9$

In order to answer this question, we will adhere to the step by step procedure given in the proof of Theorem 3.1 above. The intermediate preferences of the participants in the coalition after the first update are presented in Table 2.

Table 2: Step 1 intermediate preferences and corresponding point distribution

Preference	Fraction	A_1	A_2	A_3	A_4
$(A_2, ?, ?, ?)$	$\mathfrak{C}_{A_2} = 5/9$	0	$15/9$	0	0
(A_3, A_1, A_4, A_2)	$p_{14} = 1/9$	$2/9$	0	$3/9$	$1/9$
(A_3, A_4, A_1, A_2)	$p_{17} = 1/9$	$1/9$	0	$3/9$	$2/9$
(A_4, A_1, A_3, A_2)	$p_{20} = 2/9$	$4/9$	0	$2/9$	$6/9$
Other	$p_i = 0$	0	0	0	0
\sum	1	$7/9$	$15/9$	$8/9$	1

This allows to evaluate $d(A_2, A_i)$ for every $i \neq 2$:

$$d(A_2, A_1) = 15/9 - 7/9 = 8/9$$

$$d(A_2, A_3) = 15/9 - 8/9 = 7/9$$

$$d(A_2, A_4) = 15/9 - 1 = 6/9$$

From here we deduce that $M_1 = \max\{d(A_2, A_i) \mid i \neq k\} = d(A_2, A_1) = 8/9$ and since $w_3 \mathfrak{C}_{A_2} = 5/9 < 8/9 < 10/9 = w_2 \mathfrak{C}_{A_2}$, we are in case (b) of Step (2) above. The next step is to compute

$$t = \frac{8/9 - 1 \cdot 5/9}{2 - 1} - \varepsilon = 3/9 - \varepsilon$$

and update the intermediate profiles accordingly (see Table 3). At this point we have reduced the problem to the case of three alternatives (A_2, A_3 , and A_4) and positional rule determined by the weight vector $(3, w_{2,3}, 0)$, where

$$w_{2,3} = \frac{(2/9 + \varepsilon) \cdot 1 + (3/9 - \varepsilon) \cdot 2}{5/9} = 8/5 - 9\varepsilon/5 = 1.6 - 9\varepsilon/5.$$

Table 3: Step 2 intermediate preferences and corresponding point distribution

Preference	Fraction	A_1	A_2	A_3	A_4
$(A_2, A_1, ?, ?)$	$t = 3/9 - \varepsilon$	$6/9 - 2\varepsilon$	$1 - 3\varepsilon$	0	0
$(A_2, ?, A_1, ?)$	$2/9 + \varepsilon$	$2/9 + \varepsilon$	$6/9 + 3\varepsilon$	0	0
(A_3, A_1, A_4, A_2)	$p_{14} = 1/9$	$2/9$	0	$3/9$	$1/9$
(A_3, A_4, A_1, A_2)	$p_{17} = 1/9$	$1/9$	0	$3/9$	$2/9$
(A_4, A_1, A_3, A_2)	$p_{20} = 2/9$	$4/9$	0	$2/9$	$6/9$
Other	$p_i = 0$	0	0	0	0
\sum	1	$15/9 - \varepsilon$	$15/9$	$8/9$	1

At this stage the question of manipulability is resolved according to whether or not the inequality

$$d(A_2, A_3) + d(A_2, A_4) > w_{2,3} \mathfrak{C}_{A_2}$$

is satisfied. Plugging in the corresponding values we obtain

$$7/9 + 6/9 = 13/9 > 8/9 - \varepsilon = 5/9(1.6 - 9\varepsilon/5),$$

securing the affirmative answer.

One possible arrangement of the complete profiles is presented in Table 4 and this concludes our example.

Table 4: Complete strategic profile and resulting point distribution

Preference	Fraction	A_1	A_2	A_3	A_4
(A_2, A_1, A_3, A_4)	$(3/9 - \varepsilon) \cdot 3/5$	$18/45 - 6\varepsilon/5$	$27/45 - 9\varepsilon/5$	$9/45 - 3\varepsilon/5$	0
(A_2, A_3, A_1, A_4)	$(2/9 + \varepsilon) \cdot 3/5$	$6/45 + 3\varepsilon/5$	$18/45 + 9\varepsilon/5$	$12/45 + 6\varepsilon/5$	0
(A_2, A_1, A_4, A_3)	$(3/9 - \varepsilon) \cdot 2/5$	$12/45 - 4\varepsilon/5$	$18/45 - 6\varepsilon/5$	0	$6/45 - 2\varepsilon/5$
(A_2, A_4, A_1, A_3)	$(2/9 + \varepsilon) \cdot 2/5$	$4/45 + 2\varepsilon/5$	$12/45 + 6\varepsilon/5$	0	$8/45 + 4\varepsilon/5$
(A_3, A_1, A_4, A_2)	$p_{14} = 1/9$	$2/9$	0	$3/9$	$1/9$
(A_3, A_4, A_1, A_2)	$p_{17} = 1/9$	$1/9$	0	$3/9$	$2/9$
(A_4, A_1, A_3, A_2)	$p_{20} = 2/9$	$4/9$	0	$2/9$	$6/9$
Else	$p_i = 0$	0	0	0	0
\sum	1	$75/45 - \varepsilon$	$75/45$	$61/45 + 3\varepsilon/5$	$59/45 + 2\varepsilon/5$

Since $w_2 = w_3 = \dots = w_m = 0$ under the Plurality rule, the SI inequalities reduce to $d(A_k, A_i) > 0$ for $i \in \{1, \dots, m \mid i \neq k\}$. Then the following corollary can be deduced from Theorem 3.2.

Corollary 3.3. *Consider the Plurality voting rule with $m \geq 3$ alternatives. A profile $p = (p_1, \dots, p_{m!})$ is manipulable by a coalition with unifying alternative A_k if and only if p satisfies the following system of inequalities:*

$$\left\{ \begin{array}{l} \sum_{i=1}^{m!} p_i = 1 \\ p_i \geq 0 \\ \text{The initial arrangement is } (A_1, A_2, \dots, A_m) \\ d(A_k, A_i) > 0 \text{ for } i \in \{1, \dots, m \mid i \neq k\} \quad \text{(SI)}. \end{array} \right. \quad (3.7)$$

We would like to bring the reader's attention to the fact that the SI inequalities in (3.7) are the same as inequalities (2) in Theorem 1 of [Lepelley and Mbih \(1987\)](#), where the characterization of coalitional manipulable profiles under Plurality rule was first derived. We also bring the reader's attention to the fact that [Lepelley and Valognes \(2003\)](#) and [Favardin et al. \(2002\)](#) deal with the coalitional manipulation of the Antiplurality rule ($\lambda = 1$) and the Borda rule ($\lambda = 2$), respectively, in the case of three-alternative elections. The approach that we use in this paper to derive our list of inequalities is different from those used by [Lepelley and Valognes \(2003\)](#) and [Favardin et al. \(2002\)](#). This explains why the systems given in these papers (Lemma 2 in [Lepelley and Valognes \(2003\)](#) and Lemma 4 in [Favardin et al. \(2002\)](#)) are not a direct consequence of the systems given in Theorem 3.1. However, our results in Section 4 for the case of three alternatives allow us to show that the two characterizations are equivalent.

As already noticed, the approach that we consider in the main result of the paper can be adapted when only manipulation by small coalitions is considered. Indeed, under some

profiles a small coalition may reverse the relative ranking of two alternatives as it is shown in the following example.

Example 3.4. Let $m = 3$ and the scoring rule be Borda rule having $w = (2, 1, 0)$. As in Example 2.3, we set $p_1 = 5/9$, $p_3 = 4/9$, and $p_j = 0$ for $j \neq 1, 3$. The initial arrangement is (A_1, A_2, A_3) since alternative A_1 wins and gets $14/9$ points, A_2 gets $13/9$ points, and finally A_3 gets 0 points. But, if a fraction $p = 2/9$ of voters chose (A_2, A_3, A_1) instead of (A_2, A_1, A_3) , and the others kept their preferences unchanged, then the final result would be in favor of alternative A_2 since A_1 gets $12/9$ points, A_2 gets $13/9$ points, and A_3 gets $2/9$ points. In other words, a manipulation in favor of A_2 is also possible with a coalition of voters smaller than the maximal size (see Example 2.3 for a manipulation by a coalition of a maximal size).

Corollary 3.5. Let $m \geq 3$ and suppose that the initial arrangement is (A_1, A_2, \dots, A_m) under the positional voting rule having weight vector $w = (w_1, w_2, \dots, w_m)$. A profile $p = (p_1, \dots, p_{m!})$ is manipulable by some coalition not exceeding a proportion $0 < p \leq 1$ of the electorate with unifying alternative A_k if and only if p satisfies the following system of inequalities:

$$\left\{ \begin{array}{l} \text{Preliminary Inequalities (PI)} \\ \left\{ \begin{array}{l} \sum_{i=1}^{m!} p_i = 1 \\ p_i \geq 0 \\ 0 \leq \tilde{p}_j \leq p_j \text{ for } j, \text{ s.t. } A_1 \prec_j A_k \\ \sum_{A_1 \prec_j A_k} \tilde{p}_j = p \\ \text{The initial arrangement is } (A_1, A_2, \dots, A_m) \end{array} \right. \\ \\ \text{Strategic Inequalities (SI)} \\ \left\{ \begin{array}{l} \sum_{i \neq k} d(A_k, A_i) > (w_2 + \dots + w_{m-1} + w_m)p \\ \sum_{i \neq k} d(A_k, A_i) - M_1 > (w_3 + \dots + w_{m-1} + w_m)p \\ \dots \\ \sum_{i \neq k} d(A_k, A_i) - \sum_{j=1}^{m-4} M_j > (w_{m-2} + w_{m-1} + w_m)p \\ \sum_{i \neq k} d(A_k, A_i) - \sum_{j=1}^{m-3} M_j > (w_{m-1} + w_m)p \\ M_{m-1} > w_m p, \end{array} \right. \end{array} \right. \quad (3.8)$$

where by $A_1 \prec_j A_k$ we understand that A_k has a higher ranking than A_1 in preference of type j and \tilde{p}_j denotes the share of ‘participants’ with preference of this type. The added PI inequalities correspond to the fact that each \tilde{p}_j can not exceed p_j . As it is sufficient to consider coalitions of the largest allowed proportion, we add the equality $\sum_{A_1 \prec_j A_k} \tilde{p}_j = p$.

The next remark concerns the case where the number of voters is small.

Remark 3.6. Theorem 3.1 does not hold true in the case of finite number of voters. Consider the case of $m = 3$ alternatives A_1, A_2 and A_3 and the scoring rule with weight vector $w = (1, 0.9, 0)$. Let there be $n = 21$ voters and the profile $p = (6, 7, 8, 0, 0, 0)$. Then initially A_1 gets 20.2 points, A_2 gets 13.4 points and A_3 gets 6.3 points satisfying the arrangement assumptions. Next we show that the remaining two inequalities in the system, responsible for the possibility of manipulation by Coal_{A_2} , hold true for the profile p as well. Indeed, $d(A_2, A_1) = 0.4 > 0$ and $d(A_2, A_1) + d(A_2, A_3) = 7.5 > 7.2 = \lambda c_{A_2}$. However, if a single member of the coalition updated his/her profile to (A_2, A_1, A_3) then $\lambda = 0.9 > 0.4 = d(A_2, A_1)$ would lead to alternative A_1 getting ahead of A_2 . On the other hand it is not possible for all 8 people in Coal_{A_2} to update their profiles to (A_2, A_3, A_1) either, as $d(A_2, A_3) = 7.1 < 7.2 = \lambda c_{A_2}$. Hence, the profile is not manipulable by Coal_{A_2} .

We finish this part with a remark on the manipulability of the Antiplurality rule.

Remark 3.7. Suppose that the initial arrangement is (A_1, A_2, \dots, A_m) . The group of voters having the last alternative prior in their preferences (Coal_{A_m}) can never manipulate an election under Antiplurality rule whichever the number of alternatives m is. Otherwise, as after such a manipulation the last alternative A_m would have the same number of points (his number of points is not influenced by the manipulation) all the other alternatives would have to undergo a simultaneous loss of points. That is clearly impossible, because the number of rearranged points equals to the number of points initially arranged between the alternatives A_1, \dots, A_{m-1} .

4 Some numerical results

The aim of this section is to compute the share of manipulable profiles for three well-studied scoring rules under the assumption of Impartial and Anonymous Culture (IAC) in the limit case as $n \rightarrow \infty$. We consider this hypothesis because it has been widely used in many papers that we cite in the introduction. When $n \rightarrow \infty$, this assumption indicates that all profile vectors of the form $p = (p_1, \dots, p_{m!})$, with each $p_i \geq 0$ and $\sum_{i=1}^{m!} p_i = 1$, are equally likely to occur. This probability distribution was first introduced by [Fishburn and Gehrlein \(1976\)](#). For more details on the IAC condition and others, we refer the reader to [Diss and Merlin \(2020\)](#), [Gehrlein and Lepelley \(2011\)](#), [Gehrlein and Lepelley \(2017\)](#).

It is well-known that the limiting probability of every voting event (here the manipulation of voting outcomes) under IAC can be simply reduced to volume computations. Since the normalized profile vectors are of the form $p = (p_1, \dots, p_{m!})$ with each $p_i \geq 0$ and $\sum_{i=1}^{m!} p_i = 1$, the polytope that describes all possible profile vectors defines the standard simplex $\Delta \subset \mathbb{R}^{m!}$. The volume of $\Delta \subset \mathbb{R}^d$ is equal to $\frac{1}{(d-1)!}$, which, since in our case $d = m!$, becomes

$\frac{1}{m!-1}$. What remains to compute is the volumes of the polytopes \mathcal{P}_{A_k} (with $k = 1, \dots, m$) cutting out the regions of profiles manipulable by Coal_{A_k} inside the standard simplex Δ . For instance, in the case of three alternatives, the volumes of the polytopes \mathcal{P}_{A_2} and \mathcal{P}_{A_3} cutting out the regions of profiles manipulable by Coal_{A_2} and Coal_{A_3} (see systems in (3.1)) inside the standard simplex Δ are required. Then the total volume of the region of manipulable profiles is given by $\text{Vol}(\mathcal{P}_{A_2}) + \text{Vol}(\mathcal{P}_{A_3}) - \text{Vol}(\mathcal{P}_{A_2} \cap \mathcal{P}_{A_3})$. Afterwards the result is divided by the volume of the standard simplex and multiplied by 6 (all possible final arrangements of alternatives) to produce the final answer. The approach described in [Cervone et al. \(2005\)](#)³ tells us an effective way to tackle this problem for three-alternative elections. When we have more than three alternatives, however, those methods become too complicated in practice. Indeed, due to dimension reasons, the precise probabilities seem to be difficult to obtain for $m = 4$ and it is impossible to compute for $m \geq 5$ at this stage using the existing algorithms.

In case the precise volume cannot be found, it is natural to obtain a sufficiently good approximation of it. One of the most commonly used procedures for this purpose is the Monte Carlo volume estimation. The share of manipulable outcomes for Plurality, Antiplurality and Borda rule in the case of $m = 3, 4, 5$ alternatives are presented in [Table 5](#) (exact results), and [Tables 6](#) and [7](#) (simulated results).

Table 5: Share of manipulable outcomes with $m = 3$ alternatives

Rule	Manipulable	by Coal_{A_2}	by Coal_{A_3}
Plurality	29.17%	24.65%	15.63%
Antiplurality	51.85%	51.85%	0%
Borda	50.25%	47.71%	9.71%

Table 6: Share of manipulable outcomes with $m = 4$ alternatives

Rule	Manipulable	by Coal_{A_2}	by Coal_{A_3}	by Coal_{A_4}
Plurality	87.38%	83.65%	73.87%	63.53%
Antiplurality	87.13%	86.47%	22.83%	0%
Borda	95.65%	95.03%	79.16%	43.38%

Table 7: Share of manipulable outcomes with $m = 5$ alternatives

Rule	Manipulable	by Coal_{A_2}	by Coal_{A_3}	by Coal_{A_4}	by Coal_{A_5}
Plurality	99.51%	99.37%	99.05%	98.57%	98.04%
Antiplurality	97.15%	96.79%	54.78%	6.52%	0%
Borda	99.76%	99.23%	98.95%	98.18%	97.83%

³See also [Moyouwou and Tchantcho \(2017\)](#).

Our numerical results allow us to conclude that (i) moving from three to four and five alternatives very significantly increases the vulnerability to coalitional manipulation under all the scoring rules that we consider and (ii) the hierarchy of the scoring rules with respect to the susceptibility to coalitional manipulability changes when the number of alternatives increases: The Antiplurality rule seems to be the most performing when the number of alternatives is four and five, whereas it is the most prone to manipulation in the case of three alternatives.

Recall that [Moyouwou and Tchantcho \(2017\)](#) determined necessary and sufficient conditions for a given profile to be coalitionally manipulable under a general scoring rule with three alternatives. The share of manipulable profiles has also been given in this paper under the IAC condition. Our results for Table 5 coincide with their results which were already obtained in the literature as in [Lepelley and Mbih \(1987\)](#), [Lepelley and Mbih \(1994\)](#), and [Pritchard and Wilson \(2007a\)](#). The case $m = 4$ for the Plurality rule was studied in [El Ouafdi et al. \(2020a\)](#). The precise value of 87.28% for general manipulability was obtained (this corresponds to 87.38% appearing in the top left corner of Table 6). In other words, our approximation differs by 0.1%. Note also that the zero value in the last column of the Antiplurality rule is consistent with Remark 3.7.

Finally, the Python code for carrying out the computation of the share of manipulable outcomes in the case of Borda rule with $m = 4$ alternatives is provided in Appendix A. A detailed outline on the execution complimentary to the comments inside the code is given. The recipe for finding the corresponding results for other scoring voting rules with more than $m \geq 4$ alternatives is provided.

5 Conclusion

At least three main extensions can emerge from the present study. First, it would be interesting to investigate other voting rules not yet considered, especially in the class of the so-called *iterative scoring rules*, also known as *multi-stage* or *sequential elimination scoring rules*. This class of multi-round procedures is based on the same scoring principle previously defined but proceed by eliminating one or more alternatives at each round, until there is only one alternative left who is considered as the winner. Second, taking into account the possibility of reactions of voters outside the coalition that manipulates the vote may also be a good extension of this study. This appears to have a considerable impact on the manipulability shares and on the induced hierarchy of voting rules when analyzing strategic voting (see for instance [Favardin and Lepelley \(2006\)](#)). Third, our probabilistic investigations may also be extended to the framework with small sizes of manipulating coalitions using Corollary 3.8.

Appendix A: Python code used for computations in the case of Borda count with $m = 4$ alternatives

In this section we present the code (written in Python) used for carrying out the algorithm for finding the share of manipulable outcomes in the case of Borda rule with $m = 4$ alternatives. With a few modifications to the code provided one can perform analogous computations for scoring voting rules with different weights.

Step 1. Create the array of possible preferences with the alternatives corresponding to numbers 0 to 3 arranged in increasing order. This array has 24 elements. Each element of the preferences' array is an array of its own with 4 elements (see lines 14–25 in the code).

```
preferences[0] = {[0], [1], [2], [3]},  
preferences[1] = {[0], [1], [3], [2]},  
...  
preferences[23] = {[3], [2], [1], [0]}.
```

After this the point distribution array is created, i.e., the element (i, j) (here $0 \leq i \leq 23$ and $0 \leq j \leq 3$) is the number of points alternative j gets from preference i (lines 27–37.)

Step 2. Now we are in position to store the inequalities responsible for the initial arrangement of alternatives. As usual we assume that this arrangement is $(0, 1, 2, 3)$ and the 3 PI inequalities responsible for that are obtained (lines 3–8 and 38–41).

Step 3. Next generate a sample of n ('trials' in the code) random points in the simplex Δ . For this choose 24 random numbers on the closed interval $[0; 1]$, arrange them in a nondecreasing order and take the subsequent differences. The points created this way have a uniform distribution in the simplex Δ (see Chapter V.2 in [Devroye \(1986\)](#)). The corresponding lines of the code are 43–54.

Step 4. The coalition arrays are established. These are 3 arrays of 24 elements, each element equal to 1 or 0, depending on whether preference i belongs to the coalition or not (lines 55–66).

Step 5. The difference arrays are generated (lines 67–105).

Step 6. The remaining part of the code stores the SI inequalities (3.2) responsible for the profile being manipulable by each of the coalitions separately (recall that inequalities (3.2) are written in terms of the elements of difference arrays established on the previous step) and then checks for all sample points to determine how many of them satisfy the initial arrangement conditions and are manipulable by at least one coalition.

```

1 import random
2
3 InitArrangement = 3 * [0]
4
5 for i in range(0, 3):
6     InitArrangement[i] = []
7     for j in range(0, 24):
8         InitArrangement[i].append(0)
9 # generating array of profiles
10 preferences = []
11 for i in range(1, 25):
12     preferences.append(i)
13 counter = 0
14 for i in range(0, 4):
15     for j in range(0, 4):
16         for k in range(0, 4):
17             for l in range(0, 4):
18                 if (i != j and i != k and i != l
19                     and j != k and j != l and k != l):
20                     preferences[counter] = []
21                     preferences[counter].append(i)
22                     preferences[counter].append(j)
23                     preferences[counter].append(k)
24                     preferences[counter].append(l)
25                     counter += 1
26 # generating array of points distribution according to profiles
27 PointDistArr = []
28 for i in range(0, 4):
29     PointDistArr.append(i)
30 for i in range(0, 4):
31     PointDistArr[i] = []
32 for j in range(0, 24):
33     PointDistArr[i].append(0)
34 for i in range(0, 24):
35     for j in range(0, 4):
36         s = preferences[i][j]
37         PointDistArr[s][i] = 3 - j
38 for j in range(0, 24): # Initial arrangement is A>B>C>D
39     InitArrangement[0][j] = PointDistArr[0][j] - PointDistArr[1][j]
40     InitArrangement[1][j] = PointDistArr[1][j] - PointDistArr[2][j]
41     InitArrangement[2][j] = PointDistArr[2][j] - PointDistArr[3][j]
42
43 trials = 8000000 # number of points in the sample
44 TrialPts = trials * [0]
45 S = 24 * [0]
46 # creating random sample of points in the simplex
47 for i in range(0, trials):
48     TrialPts[i] = []
49     for j in range(0, 23):
50         S[j] = random.uniform(0, 1)
51     S[23]=1
52     TrialPts[i] = sorted(S)
53     for k in range(1, 24):

```

```

54     TrialPts[i][24 - k] -= TrialPts[i][23 - k]
55 # initiating coalition members arrays (1 if the preference
56 # belongs to the coalition and 0 otherwise)
57 CoalB = 24 * [0]
58 CoalC = 24 * [0]
59 CoalD = 24 * [0]
60 for j in range(0, 24):
61     if PointDistArr[0][j] < PointDistArr[1][j]:
62         CoalB[j] = 1
63     if PointDistArr[0][j] < PointDistArr[2][j]:
64         CoalC[j] = 1
65     if PointDistArr[0][j] < PointDistArr[3][j]:
66         CoalD[j] = 1
67 dBA = 24 * [0]
68 dBC = 24 * [0]
69 dBD = 24 * [0]
70 dCA = 24 * [0]
71 dCB = 24 * [0]
72 dCD = 24 * [0]
73 dDA = 24 * [0]
74 dDB = 24 * [0]
75 dDC = 24 * [0]
76 # computing d(B,A), d(B,C) and d(B,D)
77 for j in range(0, 24):
78     if CoalB[j] == 1:
79         dBA[j] = 3
80         dBC[j] = 3
81         dBD[j] = 3
82     else:
83         dBA[j] = PointDistArr[1][j] - PointDistArr[0][j]
84         dBC[j] = PointDistArr[1][j] - PointDistArr[2][j]
85         dBD[j] = PointDistArr[1][j] - PointDistArr[3][j]
86 # computing d(C,A), d(C,B) and d(C,D)
87 for j in range(0, 24):
88     if CoalC[j] == 1:
89         dCA[j] = 3
90         dCB[j] = 3
91         dCD[j] = 3
92     else:
93         dCA[j] = PointDistArr[2][j] - PointDistArr[0][j]
94         dCB[j] = PointDistArr[2][j] - PointDistArr[1][j]
95         dCD[j] = PointDistArr[2][j] - PointDistArr[3][j]
96 # computing d(D,A), d(D,B) and d(D,C)
97 for j in range(0, 24):
98     if CoalD[j] == 1:
99         dDA[j] = 3
100        dDB[j] = 3
101        dDC[j] = 3
102    else:
103        dDA[j] = PointDistArr[3][j] - PointDistArr[0][j]
104        dDB[j] = PointDistArr[3][j] - PointDistArr[1][j]
105        dDC[j] = PointDistArr[3][j] - PointDistArr[2][j]
106

```

```

107 counter = 0
108 B = 3*[0]
109 C = 3*[0]
110 D = 3*[0]
111 Sb = trials*[0]
112 Sc = trials*[0]
113 Sd = trials*[0]
114 BmanipCheck = False
115 CmanipCheck = False
116 DmanipCheck = False
117 CoalBsize = 0
118 CoalCsize = 0
119 CoalDsize = 0
120 for i in range(0, trials):
121     check = False
122     CoalBsize = 0
123     CoalCsize = 0
124     CoalDsize = 0
125     Sb[i] = []
126     Sc[i] = []
127     Sd[i] = []
128     for j in range(0, 3):
129         s = 0
130         check = True
131         for k in range(0, 24):
132             s += TrialPts[i][k] * InitArrangement[j][k]
133         if s < 0:
134             check = False
135             break
136     if check is True:
137         BmanipCheck = False
138         CmanipCheck = False
139         DmanipCheck = False
140         for j in range(0, 3):
141             B[j] = 0
142             C[j] = 0
143             D[j] = 0
144         for k in range(0, 24):
145             CoalBsize += TrialPts[i][k] * CoalB[k]
146             CoalCsize += TrialPts[i][k] * CoalC[k]
147             CoalDsize += TrialPts[i][k] * CoalD[k]
148             B[0] += TrialPts[i][k] * dBA[k]
149             B[1] += TrialPts[i][k] * dBC[k]
150             B[2] += TrialPts[i][k] * dBD[k]
151             C[0] += TrialPts[i][k] * dCA[k]
152             C[1] += TrialPts[i][k] * dCB[k]
153             C[2] += TrialPts[i][k] * dCD[k]
154             D[0] += TrialPts[i][k] * dDA[k]
155             D[1] += TrialPts[i][k] * dDB[k]
156             D[2] += TrialPts[i][k] * dDC[k]
157     Sb[i] = sorted(B)
158     Sc[i] = sorted(C)
159     Sd[i] = sorted(D)

```

```

160
161     if (Sb[i][0] > 0) and (Sb[i][0] + Sb[i][1] > CoalBsize) and (Sb[i][0]
162         + Sb[i][1] + Sb[i][2] > 3 * CoalBsize):
163         BmanipCheck = True
164     if (Sc[i][0] > 0) and (Sc[i][0] + Sc[i][1] > CoalCsize) and (Sc[i][0]
165         + Sc[i][1] + Sc[i][2] > 3 * CoalCsize):
166         CmanipCheck = True
167     if (Sd[i][0] > 0) and (Sd[i][0] + Sd[i][1] > CoalDsize) and (Sd[i][0]
168         + Sd[i][1] + Sd[i][2] > 3 * CoalDsize):
169         DmanipCheck = True
170
171     if check is True and (BmanipCheck is True or CmanipCheck is True or
172         DmanipCheck is True):
173         counter += 1
174
175 ans = 0.0
176 ans = float(24 * counter / 10000)
177 print(str(ans) + '%')

```

References

- Aleskerov, F. and Kurbanov, E. (1999) Degree of manipulability of social choice procedures. *Current Trends in Economics* 8:13–27.
- Chamberlin, J. (1985) An investigation into the relative manipulability of four voting systems. *Behavioral Science* 30(4):195–203.
- Cervone, D.P., Gehrlein, W.V. and Zwicker, W.S. (2005) What scoring rule maximizes Condorcet efficiency under IAC. *Theory and Decision* 58:145–185.
- Devroye, L. (1986) Nonuniform random variate generation. *Springer-Verlag, New York*
- Diss, M. (2015) Strategic manipulability of self-selective social choice rules. *Annals of Operations Research* 229:347–376.
- Diss, M. and Merlin, V. (2020) Evaluating voting systems with probability models, essays by and in honor of William Gehrlein and Dominique Lepelley. *Studies in Choice and Welfare - Springer*.
- Favardin, P. and Lepelley, D. (2006) Some further results on the manipulability of social choice rules. *Social Choice and Welfare* 26:485–509.
- Favardin, P., Lepelley, D. and Serais, J. (2002) Borda rule, Copeland method and strategic manipulation. *Review of Economic Design* 7(2):213–228.
- Fishburn, P.C. and Gehrlein, W.V. (1976) Condorcet’s paradox and anonymous preference profiles. *Public Choice* 26:1–18.
- Gehrlein, W.V. and Lepelley, D. (2011) *Voting paradoxes and group coherence*. *Studies in Choice and Welfare - Springer*.
- Gehrlein, W.V. and Lepelley, D. (2017) *Elections, Voting Rules and Paradoxical Outcomes*. *Studies in Choice and Welfare - Springer*.
- Gehrlein, W.V., Moyouwou, I. and Lepelley, D. (2013) The impact of voters’ preference diversity on the probability of some electoral outcomes. *Mathematical Social Sciences* 66:352–365.
- Gibbard, A. (1973) Manipulation of voting schemes: a general result. *Econometrica* 41:587–601.

- El Ouafdi, A., Lepelley, D. and Smaoui, H. (2020a) Probabilities of electoral outcomes: from three-alternative to four-alternative elections. *Theory and Decision* 88:205–229.
- El Ouafdi, A., Lepelley, D., Smaoui, H. and Serais, J. (2020b) Sur la manipulabilité coalitionnelle du vote par note à trois niveaux. *Working paper*.
- Kamwa, E. and Moyouwou, I. (2020) Susceptibility to manipulation by sincere truncation: the case of scoring rules and scoring runoff systems. In *Evaluating Voting Systems with Probability Models, Essays by and in honor of William Gehrlein and Dominique Lepelley*. Diss, M. and Merlin, V. (editors). Springer, Berlin.
- Kelly, J.S. (1993) Almost all social choice rules are highly manipulable, but a few aren't. *Social Choice and Welfare* 10:161–175.
- Kim, K.H. and Roush, F.W. (1996) Statistical manipulability of social choice functions. *Group Decision and Negotiation* 5:263–282.
- Lepelley, D., Louichi, A. and Smaoui, H. (2008) On Ehrhart polynomials and probability calculations in voting theory. *Social Choice and Welfare* 30(3):363–383.
- Lepelley, D. and Mbih, B. (1987) The proportion of coalitionally unstable situations under the Plurality rule. *Economics Letters* 24(4):311–315.
- Lepelley, D. and Mbih, B. (1994) The vulnerability of four social choice functions to coalitional manipulation of preferences. *Social Choice and Welfare* 11(3):253–265.
- Lepelley, D. and Valognes, F. (2003) Voting rules, manipulability and social homogeneity. *Public Choice* 116(1/2):165–184.
- Moyouwou, I. and Tchantcho, H. (2017) Asymptotic vulnerability of positional voting rules to coalitional manipulation. *Mathematical Social Sciences* 89:70–82.
- Nitzan, S. (1985) The vulnerability of point-voting schemes to preference variation and strategic manipulation. *Public Choice* 47:349–370.
- Peleg, B. (1979) A note on manipulability of large voting schemes. *Theory and Decision* 11:401–412.
- Pritchard, G. and Wilson, M. (2007a) Exact results on manipulability of positional voting rules. *Social Choice and Welfare* 29(3):487–513.
- Pritchard, G. and Wilson, M. (2007b) Probability calculations under the IAC hypothesis. *Mathematical Social Sciences* 54(3):244–256.
- Saari, D. (1990) Susceptibility to manipulation. *Public Choice* 64:21–41.
- Satterthwaite, M.A. (1975) Strategy-proofness and Arrow's conditions: Existence and correspondence theorems for voting procedures and social welfare functions. *Journal of Economic Theory* 10(2):187–217.
- Schürmann, A. (2013) Exploiting polyhedral symmetries in social choice. *Social Choice and Welfare* 40(4):1097–1110.